



Diabetes and dementia

The onset of both type 2 diabetes and dementia are strongly age-dependent. The global epidemic of type 2 diabetes and an increase in life expectancy means that the co-morbidity of diabetes and dementia is becoming increasingly common.¹ Both type 2 and type 1 diabetes are associated with cognitive decline, increasing the risk of dementia and hastening its progress compared to those without diabetes. People with dementia are likely to find managing their diabetes very challenging and, thus, may be at greater risk of diabetes complications. This *At a glance factsheet* outlines the relationships between diabetes, cognitive decline and dementia, and the consequences of dementia for the management of diabetes. An accompanying guide to managing diabetes in people with dementia [can be found here](#).

The nature and prevalence of dementia

- Dementia is a condition characterised by cognitive impairment to the degree that it compromises the ability to lead a normal life. It is typically a progressive, irreversible condition.
- Many areas of cognition can be adversely affected, including memory, attention and concentration; language; planning and organisation; problem solving and decision making; and social skills.²
- By the end of 2025, it is estimated that 1 million people in the UK will have a diagnosis of dementia, and this number is forecast to double over the next 20 years.³
- The presentation of dementia will depend on the sub-type. Alzheimer's disease is the most common form of dementia, followed by vascular dementia (*Table 1*). Other forms of dementia are Lewy body dementia and frontotemporal dementia.⁴

Table 1. Underlying causes of dementia.⁵

Condition	Underlying causes
Alzheimer's disease (~65% of cases)	Extracellular beta-amyloid plaques; intracellular neurofibrillary tangles of hyperphosphorylated tau protein
Vascular dementia (~25% of cases)	Impaired blood supply to brain; atherosclerosis
Lewy body dementia (includes Parkinson's disease dementia)	Build-up of protein deposits in the brain
Frontotemporal dementia (includes Pick's disease)	Build-up of protein deposits in the brain
Huntington's disease	Protein mutation toxic to brain cells

Diabetes as a risk factor for dementia

- Diabetes is associated with an increased risk of cognitive impairment and dementia.
 - People with type 2 diabetes have a 1.5–2.0-times greater incidence of all-cause dementia compared to the population without diabetes.⁶
 - Cognitive decline in older adults with type 2 diabetes proceeds at twice the rate of those without diabetes.⁷
 - A recent US study found the prevalence of dementia in adults with diabetes to be 13.1% in 65–74-year-olds and 24.2% in the over-75s.⁶
- The subtypes of dementia most strongly associated with type 2 diabetes are vascular dementia (2.5-fold increased risk compared to no diabetes) and Alzheimer's disease (1.5-fold increased risk).⁸
- Type 1 diabetes is also linked to a higher incidence of dementia, notably when extremes of glycaemia (low or high) have been experienced.⁹
- In the ACCORD-MIND trial in people with type 2 diabetes, higher HbA_{1c} levels were associated with lower cognitive function.¹⁰ Further studies, such as AgeCoDe, have confirmed this link.¹¹
 - However, the extension of the ACCORD-MIND trial showed that intensive glycaemic intervention did not result in improved cognitive outcomes.¹²

How does diabetes contribute to cognitive decline?

- Atherosclerotic disease of cerebral arteries contributes to cognitive decline; this pathology is more pronounced in people with diabetes.
- Vascular dementia is a relatively more common cause of dementia in people with diabetes compared to the population without diabetes.
- Chronic hyperglycaemia generates advanced glycation end-products (AGEs), with glucose covalently bonding to proteins and other important molecules, thereby damaging their function.¹³ Microvascular damage results, with accompanying inflammatory changes including acceleration of amyloid plaque formation (affecting Alzheimer's disease).
- Insulin receptors are present in the brain, notably in the cerebral cortex and the hippocampus, which is central to memory function. In people with diabetes, insulin resistance at this local level will compromise glucose utilisation, drive inflammatory change and damage neuronal function.¹⁴
- Neuronal damage can also arise in response to acute hyperglycaemia. In this situation, there can be a reduction in cerebral blood flow that exposes cells to oxidative stress and subsequent molecular damage within cells.⁹

Table 2. Predisposition of elderly people with diabetes and dementia to hypoglycaemia.

Predisposing factor	Comment
Mistakes in taking medication	Overdosing of medications that can induce hypoglycaemia
Decreased hepatic function	Reduced metabolism of antidiabetes drugs
Decreased renal function	Reduced excretion of antidiabetes drugs
Polypharmacy	Drugs inducing changes in metabolism leading to accumulation of antidiabetes drugs
Inconsistent eating patterns and missed meals	Reducing blood glucose levels

Hypoglycaemia and dementia

- There is mounting evidence that episodes of hypoglycaemia lead to cognitive decline and increase the risk of dementia. The more frequent and severe these episodes, the greater the risk.⁹
- Sulfonylureas, meglitinides and insulin are the treatments most likely to induce hypoglycaemia.
- Intensive insulin therapy, most commonly in the context of type 1 diabetes, carries the highest risk of hypoglycaemia.¹⁵
- Dementia itself increases risk of hypoglycaemia for multiple reasons (see Table 2).^{1,16} The ensuing neuroglycopenia can lead to permanent neuronal damage.⁹
- People with dementia, and elderly people in general, are at greater risk of medication-induced hypoglycaemia and of more serious consequences from this – see Table 3.

Depression, diabetes and dementia

- Depression is a risk factor for dementia.
- People with type 2 diabetes are more prone to depression than those without diabetes.⁶
- Individuals with diabetes who suffer from depression are 2.7-times more likely to develop dementia than those with diabetes alone.¹⁷

Table 3. Problems associated with medication-induced hypoglycaemia in elderly people with diabetes and dementia.

Problem	Comment
Reduced hypoglycaemic awareness	Greater risk of neuroglycopenia
Inadequate physiological response to hypoglycaemia	Due to reduced counter-regulatory hormones (principally glucagon and adrenaline, which stimulate release of glucose from the liver)
Reduced ability to manage episodes of hypoglycaemia	Consequence of dementia and frailty
More severe consequences of hypoglycaemia	Examples include falls, with risk of subdural haemorrhage and osteoporotic fracture; risk of cardiac arrhythmias and cardiovascular events

Consequences of dementia for managing diabetes

- Education and empowerment of an individual to self-manage their diabetes is compromised with the co-morbidity of dementia.¹⁸
- People with dementia may lack the ability to make healthy lifestyle choices that are important in managing their diabetes, notably diet and exercise.¹⁹
- Medications may be taken incorrectly or not at all.
 - In the case of antidiabetes medications, this places the individual at risk of both hyper- and hypoglycaemia, both of which can accelerate cognitive decline.¹⁶
 - Missing treatments for hypertension and dyslipidaemia increases the risk of cardiovascular complications.
- Self-monitoring of blood glucose levels may not be possible, so that persistent hyperglycaemia or episodes of hypoglycaemia pass unidentified.
- In addition to problems with memory, organisation and planning, people with dementia may be further burdened with behavioural and psychological symptoms that compromise diabetes management.²⁰



Key learning points

- With an ageing population, the incidence of both dementia and diabetes is rising and the co-morbidity of these two conditions is increasingly common. This has profound personal and social consequences.
- People with diabetes have an increased risk of cognitive decline and dementia. Type 2 diabetes is associated with a 2.5-fold increased risk of vascular dementia and a 1.5-fold increased risk of Alzheimer's disease.
- Both hyperglycaemia and medication-induced hypoglycaemia are responsible for cognitive decline in people with diabetes.
- People with dementia are likely to have difficulty managing their diabetes, including making healthy lifestyle choices, adhering correctly to medication regimens and monitoring blood glucose levels.

Part 2

How to approach and manage diabetes in people with dementia

[Click here to access](#)

Useful resources

- [How to manage diabetes in later life](#)
- [How to prevent, identify and manage hypoglycaemia in adults with diabetes](#)
- [At a glance factsheet: Deprescribing in type 2 diabetes](#)
- [Diabetes Distilled: The 4S Pathway – realigning management for older people with diabetes](#)

References

1. Bunn F, Goodman C, Jones PR et al (2017) Managing diabetes in people with dementia: A realist review. *Health Technol Assess* **21**: 1–140
2. NHS (2024) *What is dementia?* Available at: <http://bit.ly/4k93ZX9>
3. Dementia UK (2023) *What is dementia?* Available at: <http://bit.ly/4njhFSa>
4. Larson ST, Swegle J, Fishbein E, Wilbur J (2023) Dementia: dementia types. *FP Essentials* **534**: 7–11
5. Kuate Defo A, Bakula V, Pisaturo A et al (2024) Diabetes, antidiabetic medications and risk of dementia: A systematic umbrella review and meta-analysis. *Diabetes Obes Metab* **26**: 441–62
6. Reddy S, Hashmi A (2020) Managing diabetes and dementia. *Clin Geriatr Med* **36**: 419–29
7. Cholerton B, Baker LD, Montine TJ, Craft S (2016) Type 2 diabetes, cognition, and dementia in older adults: Toward a precision health approach. *Diabetes Spectr* **29**: 210–9
8. Cheng G, Huang C, Deng H, Wang H (2012) Diabetes as a risk factor for dementia and mild cognitive impairment: A meta-analysis of longitudinal studies. *Intern Med J* **42**: 484–91
9. Husain KH, Sarhan SF, AlKhalifa HKAA et al (2023) Dementia in diabetes: The role of hypoglycemia. *Int J Mol Sci* **24**: 9846
10. Cukierman-Yaffe T, Gerstein HC, Williamson JD et al; ACCORD-MIND investigators (2009) Relationship between baseline glycemic control and cognitive function in individuals with type 2 diabetes and other cardiovascular risk factors: The Action to Control Cardiovascular Risk in Diabetes–Memory in Diabetes (ACCORD-MIND) trial. *Diabetes Care* **32**: 221–6
11. Ramirez A, Wolfgruber S, Lange C et al; AgeCoDe Study Group (2015) Elevated HbA_{1c} is associated with increased risk of incident dementia in primary care patients. *J Alzheimers Dis* **44**: 1203–12
12. Murray AM, Hsu FC, Williamson JD et al; ACCORDION-MIND investigators (2017) ACCORDION-MIND: results of the observational extension of the ACCORD-MIND randomised trial. *Diabetologia* **60**: 69–80
13. Singh VP, Bali A, Singh N, Jaggi AS (2014) Advanced glycation end products and diabetic complications. *Korean J Physiol Pharmacol* **18**: 1–14
14. Chen W, Cai W, Hoover B, Kahn CR (2022) Insulin action in the brain: Cell types, circuits, and diseases. *Trends Neurosci* **45**: 384–400
15. Nakhleh A, Shehadeh N (2021) Hypoglycemia in diabetes: An update on pathophysiology, treatment, and prevention. *World J Diabetes* **12**: 2036–49
16. Punthakee Z, Miller ME, Launer LJ et al; ACCORD investigators; ACCORD-MIND investigators (2012) Poor cognitive function and risk of severe hypoglycemia in type 2 diabetes: *Post hoc* epidemiologic analysis of the ACCORD trial. *Diabetes Care* **35**: 787–93
17. Aderinto N, Olatunji G, Abdulbasit M et al (2023) The impact of diabetes in cognitive impairment: A review of current evidence and prospects for future investigations. *Medicine (Baltimore)* **102**: e35557
18. Sinclair AJ, Girling AJ, Bayer AJ (2000) Cognitive dysfunction in older subjects with diabetes mellitus: Impact on diabetes self-management and use of care services. All Wales Research into Elderly (AWARE) Study. *Diabetes Res Clin Pract* **50**: 203–12
19. Feil DG, Zhu CW, Sultzer DL (2012) The relationship between cognitive impairment and diabetes self-management in a population-based community sample of older adults with type 2 diabetes. *J Behav Med* **35**: 190–9
20. Tomlin A, Sinclair A (2016) The influence of cognition on self-management of type 2 diabetes in older people. *Psychol Res Behav Manag* **9**: 7–20

Author: David Morris, retired GP and Specialist Doctor in Diabetes; Undergraduate Clinical Tutor, Keele University.

Citation: Morris D (2025) At a glance factsheet: Diabetes and dementia. *Diabetes & Primary Care* **27**: 111–3