Clarke score⁷

Answers provide a rating of either A (aware) or R (reduced).

Four or more R ratings suggest impaired hypoglycaemia awareness; ≤2=normal awareness; 3=borderline.

Question	Rating (A or R)
1. Check the category that best describes you (check one only):	
\Box I always have symptoms when my blood sugar is low (A)	
 I sometimes have symptoms when my blood sugar is low (R) I no longer have symptoms when my blood sugar is low (R) 	
 2. Have you lost some of the symptoms that used to occur when your blood sugar was low? ❑ Yes (R) ❑ No (A) 	
3. In the past six months , how often have you had moderate hypoglycaemia episodes? (episodes where you might feel confused, disoriented, or lethargic and were unable to treat yourself)	
$\square Never (A) \square Once or twice (R) \square Every other month (R) \square Once a month (R) \square More than once a month (R)$	
4. In the past year , how often have you had severe hypoglycaemic episodes? (episodes where you were unconscious or had a seizure and needed glucagon or intravenous glucose)	
(A if never; R if one or more. If ≥12, conclude impaired awareness irrespective of other answers)	
 5. How often in the last month have you had blood sugar readings under 3.9 mmol/L with symptoms? Never 1 1–3 times 1 per week 2–3 per week 2 >4–5 per week 2 >Almost daily (see question 6 for scoring) 	
 6. How often in the last month have you had blood sugar readings under 3.9 mmol/L without symptoms? □ Never □ 1–3 times □ 1 per week □ 2–3 per week □ >4–5 per week □ >Almost daily (R = answer to Q6 < answer to Q6 < answer to Q5) 	
7. How low does your blood sugar need to go before you feel symptoms? 3.4–3.8 mmol/L (A) 2.8–3.3 mmol/L (A) 2.2–2.7 mmol/L (R) Under 2.2 mmol/L (R)	
8. To what extent can you tell by your symptoms that your blood sugar is low? □ Never (R) □ Rarely (R) □ Sometimes (R) □ Often (A) □ Always (A)	