

# An exploration of the relationship between quality of life, emotional distress and HbA<sub>1c</sub> in paediatric diabetes service users

Maria Langridge, Jenny Cropper, Sujata Edate

**The prevalence of both type 1 diabetes and type 2 diabetes in children and young people is continuing to increase across most populations globally. As the prevalence of diabetes increases in this cohort, the burden of the condition will also increase. This paper discusses the role of two psychosocial correlates, emotional distress and quality of life, in the context of HbA<sub>1c</sub> outcomes in a sample of paediatric service users with diabetes. As part of this analysis, the role of age and gender are considered in the context of HbA<sub>1c</sub> outcomes. For further understanding of where resources need to be targeted, relationships between individual factors are explored. Based on our findings, we make several recommendations, but ultimately it was determined there is a need for a more holistic, person-centred approach in paediatric diabetes management.**

Psychological wellbeing constructs, such as positive affect, have been associated with better diabetes outcomes, especially glycaemic control (Massey et al, 2019). Ruiz-Aranda et al (2018) suggest that emotional distress, another psychological construct, plays a role in diabetes outcomes by influencing the acquisition and maintenance of health habits correlated with HbA<sub>1c</sub> outcomes.

Emotional distress has previously been correlated with quality of life (QoL) by Faridah et al (2017). QoL is a subjective concept of satisfaction with life and an overall sense of health and wellbeing. QoL can powerfully predict engagement with diabetes management, and thus influence long-term health outcomes (Rubin, 2000).

Identifying the factors involved in influencing HbA<sub>1c</sub> control will contribute towards the facilitation of effective and targeted interventions, which will improve the psychosocial health of children and young people (CYP) with diabetes (Hsu et al, 2018).

## Materials and procedure

A total of 190 screening questionnaires were administered to CYP with diabetes, aged 8–16 years (mean, 13 years; standard deviation [SD], 2.4), as part of their routine annual review clinic. CYP attend one annual review appointment per year, as part of their diabetes care.

The sample was 52.6% male ( $n=100$ ); most of our study sample (96.3%) had a diagnosis of type 1 diabetes (T1D;  $n=183$ ). Type 2 diabetes (T2D;  $n=5$ ) made up just 2.6% of the sample and maturity-onset diabetes of the young (MODY;  $n=2$ ) comprised 1.1% of the sample. Data were collected between January 2017 and October 2019; some CYP are represented more than once in the data due to attending multiple annual review appointments during the study period (approximately 3 years).

The clinics were held in a district general hospital where CYP received their diabetes care; the sample is therefore opportunistic, comprising only CYP attending the clinic.

**Citation:** Langridge M, Cropper J, Edate S (2022) An exploration of the relationship between quality of life, emotional distress and HbA<sub>1c</sub> in paediatric diabetes service users. *Diabetes Care for Children & Young People* 11: DCCYP092

## Article points

1. The importance of integrating psychological intervention into diabetes care has been highlighted in numerous studies.
2. The young person should be viewed holistically, beyond sole considerations of biological markers.
3. Quality of life and emotional distress are important outcomes to measure in diabetes.
4. Integrating psychology into the multidisciplinary team facilitates more informative and supportive conversations with young people and their families.
5. This study has found that concepts of emotional wellbeing and quality of life are correlated with glycaemic control.

## Key words

- Emotional distress
- HbA<sub>1c</sub>
- Holistic approach
- Psychology
- Quality of life

## Authors

Authors' details can be found at the end of the article.

**Page points**

1. The Paediatric Index of Emotional Distress was used to provide a reliable and valid measure of emotional distress in CYP aged 8–16 years, by measuring anxiety and depression within a paediatric clinical health setting.
2. The Paediatric Quality of Life Inventory was used as a self-report screening tool for quality of life scores.
3. Statistical analyses were carried out on the data to determine the strength and direction of the relationships between quality of life, emotional distress and HbA<sub>1c</sub>.

The Paediatric Index of Emotional Distress (PI-ED; O'Connor et al, 2016) was used to provide a reliable and valid measure of emotional distress in CYP aged 8–16 years, by measuring anxiety and depression within a paediatric setting. The PI-ED is a 14-item questionnaire that scores using a 4-point Likert scale (3=always, 2=a lot of the time, 1=sometimes, 0=not at all). Higher scores indicate higher levels of emotional distress; the PI-ED clinical threshold is 20.

The Paediatric Quality of Life Inventory (PedsQL Diabetes Module Version 3.0) is a brief, self-report screening tool for QoL. It consists of 28 items and five dimensions: Diabetes (11 items), Treatment I (4 items), Treatment II (7 items), Worry (3 items), and Communication (3 items). Each item is scored on a 5-point Likert scale (0=never, 1=almost never, 2=sometimes, 3=often, 4=almost always). The measures used were self-reported to eliminate the potential bias introduced by parent-proxy reports.

Blood analysis, via a Siemens DCA Vantage Analyser®, was conducted at each annual review to acquire an accurate HbA<sub>1c</sub> reading.

Statistical analyses were carried out on the

data to determine the strength and direction of the relationships between QoL, emotional distress and HbA<sub>1c</sub>. Incomplete data sets were removed to maintain the psychometric integrity of the findings.

**Results**

Statistical analyses were used to examine the relationship between QoL (mean=324.82, SD=69.54), emotional distress (mean=10.69, SD=6.81), and HbA<sub>1c</sub> control (mean=64.11 mmol/mol, SD=14.28). The mean HbA<sub>1c</sub> level of this cohort of young people was above the target 48 mmol/mol, as recommended by NICE guidelines. Levels of emotional distress, as indicated by the mean PI-ED outcome, were below the clinical threshold. Moreover, higher scores on the PedsQL indicate a better QoL, therefore the scores indicate relatively good QoL.

Using a bivariate correlation, it was found that female participants had a higher mean HbA<sub>1c</sub> level (mean=66.03 mmol/mol, SD=14.73) than males (mean=62.37 mmol/mol, SD=13.71); however, the relationship was not statistically significant ( $r=-0.107$ ;  $P=0.141$ ). Females also




Figure 1. A visual depiction of the relationship between HbA<sub>1c</sub> levels and emotional distress scores in a population of paediatric diabetes service users.

reported higher levels of emotional distress (mean=11.14, SD=7.29) than male participants (mean=10.29, SD=6.37). In continuation, females reported a lower mean QoL (mean=313.12, SD=78.16) than males (mean=335.34, SD=59.19).

Furthermore, the data indicated a statistically significant positive correlation between age and HbA<sub>1c</sub> levels ( $r=0.18$ ;  $P=0.016$ ), suggesting that older CYP have higher HbA<sub>1c</sub> levels. This is supported by the outcomes demonstrating that 16-year-old service users had the highest mean HbA<sub>1c</sub> (mean=69.04 mmol/mol, SD=16.75), whereas 9-year-old children had the lowest (mean=56.1 mmol/mol, SD=8.95).

The Pearson correlation coefficient for HbA<sub>1c</sub> and emotional distress depicts a statistically significant, positive linear relationship ( $r=0.15$ ;  $P=0.02$ ), suggesting that those experiencing lower levels of emotional distress have lower HbA<sub>1c</sub> levels (Figure 1). As emotional distress scores increase, HbA<sub>1c</sub> levels also increase.

HbA<sub>1c</sub> and mean QoL scores also demonstrate a statistically significant relationship; however, it is a moderate, negative linear relationship ( $r=-0.13$ ;  $P=0.04$ ), suggesting that those with higher HbA<sub>1c</sub> levels have lower mean QoL scores (Figure 2). As HbA<sub>1c</sub> decreases, mean QoL scores increase.

Moreover, emotional distress and QoL demonstrated a statistically significant, moderate, negative linear relationship ( $r=-0.64$ ;  $P<0.001$ ), suggesting those with higher mean QoL scores experience lower levels of emotional distress.

## Discussion

Despite advances in pharmacological treatment and intervention delivery, diabetes management remains a challenge for many CYP. The effects of diabetes and its management extend beyond purely medical outcomes. Suboptimal glycaemic control in CYP with diabetes is hypothesised to be associated with a lower reported QoL (Hassan et al, 2006), and an increased risk of emotional distress (Jones et al, 2015).

Considering the above, the aim of the present study was to correlate concepts of QoL and emotional distress with the clinical pictures of a sample of CYP with diabetes; it was hoped that this knowledge could facilitate the implementation of targeted and effective




Figure 2. A visual depiction of the relationship between HbA<sub>1c</sub> levels and quality of life scores in a population of paediatric diabetes service users.

multidisciplinary team (MDT) intervention, as well as to inform other clinicians working in paediatric diabetes teams.

The mean HbA<sub>1c</sub> in this cohort of CYP was above the NICE-recommended threshold of 48 mmol/mol. Moreover, a significant positive correlation between age and HbA<sub>1c</sub> level was found, indicating declining HbA<sub>1c</sub> control as CYP develop. This relationship can be explained, in part, in the context of age and independence; the current sample's mean age was 13 years old, the beginning of adolescence and independence. Parents and carers hold more responsibility for condition management in younger children, while adolescents are more independent (Nardi et al, 2008). Condition management is challenging and it is reported that only 20% of adolescents adhere fully to recommended condition management (Kyngäs, 2000).

It is important to acknowledge that children face multiple transitions as they age, including puberty, which can be emotionally distressing (Byrne et al, 2007). Future research should explore gender differences in the context of puberty and hormones, and how these correlate to emotional distress and QoL outcomes. This analysis would allow brief advice and information provided by medical teams to be individualised, more idealised, and attuned to the service user's needs.

### Page points

1. The mean Paediatric Index of Emotional Distress score reported by our study sample was below the recommended clinical threshold. This may be explained, in part, by the existing role of clinical psychology in the team.
2. The present study findings also highlighted that young people with a higher quality of life report lower levels of emotional distress, which is consistent with other literature.
3. Data from this study indicate that young people with higher HbA<sub>1c</sub> levels report lower quality of life. This could be explained by increased frequency of appointments and condition management causing disruption to daily routines.

While the clinical outcomes of diabetes, such as HbA<sub>1c</sub>, are fundamental aspects of diabetes care, paediatric diabetes teams should adopt a holistic model of care (Goss, et al, 2010). In agreement, our data identified higher reported emotional distress in those with higher HbA<sub>1c</sub> levels. These findings emphasise the importance of taking into consideration the emotional distress correlated with the clinical outcomes of young people with diabetes; it may worsen and result in poor condition management (Nardi et al, 2008). Given the role of emotional distress in paediatric diabetes, clinical psychologists have a key role in maximising the benefits of treatment; facilitating a safe and non-judgemental space for the acknowledgement and processing of affect.

Interestingly, the mean PI-ED score reported by our study sample was below the recommended clinical threshold. This may be explained, in part, by the existing MDT in place in the clinic, which has embraced the role of clinical psychology as an essential part of effective diabetes intervention. Moreover, much of the existing literature exploring emotional distress outcomes in CYP use outcome measures containing somatic items, which are at risk of inflation when completed by CYP with chronic illness (O'Connor et al, 2016). Our use of the PI-ED eliminated this risk, thus potentially offering an explanation for the lower reported emotional distress in our study.

With these outcomes in mind, a fundamental role of the MDT is to consider the weighting of emotional distress in the management of diabetes in CYP. The source and severity of emotional distress should be identified and measured to allow accurate formulations and, in turn, more targeted patient-centred interventions.

A study by Babenko et al (2019) highlighted that QoL is highly impacted in individuals with diabetes. Interestingly, the current sample reported higher QoL scores, indicating an overall satisfactory QoL. As described with emotional distress, this outcome may be explained by the existing MDT support, which is inclusive of clinical psychology, within in the clinic.

The present study findings also highlighted that young people with a higher QoL report lower levels of emotional distress. The consistency of such findings in current literature, in conjunction

with our data, demonstrates a relationship between diabetes, emotional distress and QoL. Therefore, diabetes interventions should acknowledge the impact of emotional wellbeing and QoL on this cohort, and integrate approaches targeted to improve these outcomes (Schram et al, 2009).

Additionally, our data indicate that CYP with higher HbA<sub>1c</sub> levels report lower QoL. Our findings can, in part, be explained by inconvenience resulting from increased frequency of appointments, as well as condition management causing disruption to daily routines (Rubin, 2000). This outcome demonstrates a pressing need for access to effective clinic-based interventions to improve QoL and HbA<sub>1c</sub> outcomes, and justifies our efforts to assess QoL perception in this cohort (Christie et al, 2014). If having a high HbA<sub>1c</sub> is correlated with a lower QoL, then this is another reason to consider and measure QoL in a clinic population.

While it is difficult to establish a causal relationship from correlational data, the relationship between QoL and HbA<sub>1c</sub> is so consistent that it is appropriate to assume that the relationship between metabolic control and QoL is equally important. Our findings validate the importance of monitoring QoL in this cohort, as it may uncover obstacles to effective condition management, and contribute to the attainment of treatment goals. Consultation with mental health professionals is invaluable in assisting diabetes care providers in the scoring and interpretation of QoL screening measures, and decision making around referrals for more in-depth psychological support (Hilliard et al, 2018).

In line with our findings and ISPAD recommendations (Delamater et al, 2018), resources should be made available to include psychologists in the MDT, who specialise in the psychological and behavioural functioning of CYP with chronic health conditions. They should be available to interact with CYP at outpatient clinics to conduct assessment of psychosocial functioning and psychological wellbeing, as well as to support the diabetes team in the identification and management of psychological and behavioural difficulty. Screening for

emotional distress and QoL should be routinely carried out, and target those with HbA<sub>1c</sub> levels outside of the recommended clinical threshold. It is imperative to identify emotional distress and QoL concerns in CYP with diabetes, as these may act as a barrier to the coordination of effective treatment for diabetes.

Future research should include a larger sample size, particularly of children with type 2 diabetes, to help identify whether there are any significant differences in emotional distress and QoL outcomes. This will further allow the MDT to tailor interventions based on individual need.

## Conclusion

The current study has contributed to the literature by providing a more comprehensive picture of health and wellbeing in CYP with diabetes. The data support the value of attending to psychological wellbeing as worthy of individual attention in CYP with diabetes, and should not be contingent upon the sole consideration of physiological markers. Clinical, personal and psychological characteristics appear to be correlated with HbA<sub>1c</sub> control, and their identification is crucial in addressing strategies to optimise HbA<sub>1c</sub> control.

A routine assessment of emotional wellbeing and QoL facilitates the identification of these factors. Having a psychologist integrated into the MDT can address the emotional and QoL-related needs of CYP with diabetes. The current findings can assist the MDT in directing resources towards those CYP who require it the most. ■

Babenko AY, Mosikian AA, Lebedev DL et al (2019) Mental state, psychoemotional status, quality of life and treatment compliance in patients with type 2 diabetes mellitus. *J Comp Eff Res* **8**: 113–20

- Byrne DG, Davenport SC, Mazanov J (2007) Profiles of adolescent stress: The development of the adolescent questionnaire (ASQ). *J Adolesc* **30**: 393–416
- Christie D, Thompson R, Sawtell M et al (2014) Structured, intensive education maximising engagement, motivation and long-term change for children and young people with diabetes: a cluster randomised controlled trial with integral process and economic evaluation—the CASCADE study. *Health Technol Assess* **18**: 1–202
- Delamater AM, de Wit M, McDarby V et al (2018) ISPAD Clinical Practice Consensus Guidelines 2018: Psychological care of children and adolescents with type 1 diabetes. *Pediatr Diabetes* **19**: Suppl 27: 237–49
- Faridah IN, Perwitasari DA, Pusita M, Jasma H (2017) Relationship between emotional distress and quality of life on type 2 diabetes mellitus patients in Meranti island regency hospital. *IOP Conf Ser: Mater Sci Eng* **259**: 012002
- Goss PW, Paterson MA, Renalson J (2010) A ‘radical’ new rural model for pediatric diabetes care. *Pediatr diabetes* **11**: 296–304
- Hassan K, Loar R, Anderson BJ, Heptulla RA (2006) The role of socioeconomic status, depression, quality of life, and glycemic control in type 1 diabetes mellitus. *J Pediatr* **149**: 526–31
- Hilliard ME, De Wit M, Wasserman RM et al (2018) Screening and support for emotional burdens of youth with type 1 diabetes: Strategies for diabetes care providers. *Pediatr Diabetes* **19**: 534–43
- Hsu HC, Lee YJ, Wang RH (2018) Influencing pathways to quality of life and HbA<sub>1c</sub> in patients with diabetes: A longitudinal study that inform evidence-based practice. *Worldviews Evid Based Nurs* **15**: 104–12
- Jones A, Vallis M, Pouwer F (2015) If it does not significantly change HbA<sub>1c</sub> levels why should we waste time on it? A plea for the prioritization of psychological well-being in people with diabetes. *Diabet Med* **32**: 155–63
- Kyngäs H (2000) Compliance of adolescents with diabetes. *J Pediatr Nurs* **15**: 260–7
- Massey CN, Feig EH, Duque-Serrano L et al (2019) Well-being interventions for individuals with diabetes: a systematic review. *Diabetes Res Clin Pract* **147**: 118–33
- Nardi L, Zucchini S, D’Alberton F et al (2008) Quality of life, psychological adjustment and metabolic control in youths with type 1 diabetes: A study with self- and parent-report questionnaires. *Pediatr Diabetes* **9**: 496–503
- O’Connor S, Ferguson E, Carney T et al (2016) The development and evaluation of the paediatric index of emotional distress (PI-ED). *Soc Psychiatry Psychiatr Epidemiol* **51**: 15–26
- Rubin RR (2000) Diabetes and quality of life. *Diabetes Spectr* **13**: 21
- Ruiz-Aranda D, Zysberg L, García-Linares E et al (2018) Emotional abilities and HbA<sub>1c</sub> levels in patients with type 1 diabetes. *Psychoneuroendocrinology* **93**: 118–23
- Schram MT, Baan CA, Pouwer F (2009) Depression and quality of life in patients with diabetes: a systematic review from the European depression in diabetes (EDID) research consortium. *Curr Diabetes Rev* **5**: 112–9

**“Having a psychologist integrated into the MDT can address the emotional and QoL-related needs of CYP with diabetes.”**

## Authors

Maria Langridge is Assistant Psychologist, Dental and Oral-maxillofacial Trauma Psychology (DOT-P) Service, North East London Foundation Trust; Jenny Cropper is Consultant Clinical Psychologist, Spires Cleft Centre & Oxford Craniofacial Unit, John Radcliffe Hospital, Oxford; Sujata Edate is Consultant Paediatrician, Paediatric Diabetes Team, Wexham Park Hospital, Frimley Health NHS Foundation Trust.