



## SGLT2 inhibitors: Indications, doses and licences

### Indications, doses and licences of SGLT2 inhibitors, by indication.

Indication	Drug and dose	Initiate	Stop/reduce	Notes
Glycaemic control	<b>Canagliflozin 100 mg</b> Increase to <b>300 mg</b> if required	eGFR $\geq$ 45 eGFR $\geq$ 60	eGFR persistently $<$ 45 unless ACR $>$ 30 Reduce to 100 mg if eGFR $<$ 60	Due to their mode of action, all four SGLT2 inhibitors have reduced glucose-lowering effects at eGFR $<$ 45. Follow the guidance on when to stop each drug and consider adding in another glucose-lowering drug if HbA <sub>1c</sub> is above the agreed, individualised, target
	<b>Dapagliflozin 10 mg</b>	eGFR $>$ 15	eGFR $<$ 15	
	<b>Empagliflozin 10 mg</b> Increase to <b>25 mg</b> if required	eGFR $\geq$ 60 eGFR $\geq$ 60	eGFR persistently $<$ 45 Reduce to 10 mg if eGFR $<$ 60	
	<b>Ertugliflozin 5 mg</b> Increase to <b>15 mg</b> if required	eGFR $\geq$ 60 eGFR $\geq$ 60	eGFR persistently $<$ 45 eGFR persistently $<$ 45	
Chronic kidney disease (CKD)	<b>Dapagliflozin 10 mg</b>	eGFR $\geq$ 15	eGFR $<$ 15	Use with other <b>CKD</b> therapies. With or without type 2 diabetes. Not recommended to initiate if eGFR $<$ 15 as no clinical experience
Diabetic kidney disease (DKD)	<b>Canagliflozin 100 mg</b>	eGFR $\geq$ 45 eGFR $\geq$ 30 <b>and</b> ACR $\geq$ 30	eGFR persistently $<$ 45 and ACR $<$ 30 Can continue to dialysis/transplant if ACR $\geq$ 30	Add on to standard of care (e.g. ACEi or ARB) for <b>DKD</b>
Symptomatic chronic HFrEF	<b>Dapagliflozin 10 mg</b>	eGFR $\geq$ 15	eGFR $<$ 15	With or without type 2 diabetes
	<b>Empagliflozin 10 mg</b>	eGFR $\geq$ 20	eGFR $<$ 20	With or without type 2 diabetes

eGFR presented in mL/min/1.73 m<sup>2</sup>; ACR presented in mg/mmol.

ACEi=angiotensin-converting enzyme inhibitor; ACR=albumin:creatinine ratio; ARB=angiotensin receptor blocker; eGFR=estimated glomerular filtration rate; HFrEF=heart failure with reduced ejection fraction.

Information correct on 5<sup>th</sup> November 2021.

Always consult the electronic BNF or the Summaries of Product Characteristics (SPCs) prior to prescribing any drug.

SPCs: [Canagliflozin](#) | [Dapagliflozin](#) | [Empagliflozin](#) | [Ertugliflozin](#)

**Author:** Pam Brown, GP, Swansea

**Citation:** Brown P (2021) SGLT2 inhibitors: Indications, doses and starting/stopping recommendations. *Diabetes & Primary Care* 23: [early view publication]



Indications, doses and starting/stopping recommendations of SGLT2 inhibitors, by drug name.

Drug	Indication	Drug and dose	Initiate	Stop/reduce	Notes
Canagliflozin	Glycaemic control	Canagliflozin 100 mg Increase to 300 mg if required	eGFR ≥45 eGFR ≥60	eGFR persistently <45 unless ACR >30 Reduce to 100 mg if eGFR <60	eGFR <45, reduced glucose-lowering effect; add another glucose-lowering drug if needed
	Diabetic kidney disease	Canagliflozin 100 mg	eGFR ≥45 eGFR ≥30 and ACR ≥30	eGFR persistently <45 and ACR <30 Can continue to dialysis/transplant if ACR ≥30	Add on to standard of care (e.g. ACEi or ARB) for <b>diabetic kidney disease</b>
Dapagliflozin	Glycaemic control	Dapagliflozin 10 mg	eGFR >15	eGFR <15	eGFR <45, reduced glucose-lowering effect; add another glucose-lowering drug if needed
	Chronic kidney disease	Dapagliflozin 10 mg	eGFR ≥15	eGFR <15	Use with other <b>CKD</b> therapies. With or without type 2 diabetes. Not recommended to initiate if eGFR <15 as no clinical experience
	Symptomatic chronic HFrEF	Dapagliflozin 10 mg	eGFR ≥15	eGFR <15	With or without type 2 diabetes
Empagliflozin	Glycaemic control	Empagliflozin 10 mg Increase to 25 mg if required	eGFR ≥60 eGFR ≥60	eGFR persistently <45 Reduce to 10 mg if eGFR <60	eGFR <45, reduced glucose-lowering effect; add another glucose-lowering drug if needed
	Symptomatic chronic HFrEF	Empagliflozin 10 mg	eGFR ≥20	eGFR <20	With or without type 2 diabetes
Ertugliflozin	Glycaemic control	Ertugliflozin 5 mg Increase to 15 mg if required	eGFR ≥60 eGFR ≥60	eGFR persistently <45 eGFR persistently <45	eGFR <45, reduced glucose-lowering effect; add another glucose-lowering drug if needed

eGFR presented in mL/min/1.73 m<sup>2</sup>; ACR presented in mg/mmol.

ACEi=angiotensin-converting enzyme inhibitor; ACR=albumin:creatinine ratio; ARB=angiotensin receptor blocker; eGFR=estimated glomerular filtration rate; HFrEF=heart failure with reduced ejection fraction.

Information correct on 5<sup>th</sup> November 2021.

Always consult the electronic BNF or the Summaries of Product Characteristics (SPCs) prior to prescribing any drug.

SPCs: [Canagliflozin](#) | [Dapagliflozin](#) | [Empagliflozin](#) | [Ertugliflozin](#)