

Type 2 diabetes rehabilitation: an exercise and education programme

Samantha Whittle, Olivia Bailey, Helen Partridge and Janet James

Structured education is central to supporting type 2 diabetes self-management, but sustaining lifestyle change is challenging. Refocusise integrates an established diabetes self-management education curriculum with supervised exercise in an 8-week programme delivered in routine care. Across 12 courses ($n=149$; mean age 64 years), participants completed pre- and post-programme assessments (including HbA_{1c}, blood pressure, BMI, lipids and HADS). Post-programme outcomes showed reductions in HbA_{1c} (-9 mmol/mol), total cholesterol (-0.3 mmol/L), systolic/diastolic BP ($-4.2/-3.8$ mmHg), BMI (-0.6 kg/m²) and HADS-D (-1.6). Improvements in HbA_{1c} were maintained at 1 year ($n=73$; -11 mmol/mol vs baseline) and 2 years ($n=27$). This service evaluation suggests that combining structured exercise with diabetes education can deliver meaningful and sustained improvements, supporting wider implementation within routine diabetes care.

Type 2 diabetes currently affects 7% of the total population in the UK, with a new diagnosis made every two minutes (Whicher et al, 2020). The prevalence has doubled over the past 20 years, with 4.7 million people now affected. This figure is projected to rise to 5.5 million by 2030, equating to around one in ten adults (Diabetes UK, 2024). The economic burden of diabetes is substantial, accounting for approximately 10% of total NHS expenditure, highlighting the need for cost-effective interventions to improve glycaemic control and reduce complications (Hex et al, 2012).

The long-term effects of type 2 diabetes are well documented and include microvascular complications, such as retinopathy, neuropathy and nephropathy, as well as macrovascular disease (Fowler, 2008). The condition is thought to contribute to 530 myocardial infarctions each week in the UK (NHS Digital, 2016), and approximately 18% of hospital beds are occupied

by patients with diabetes (NHS Digital, 2020). Hospital stays tend to be longer for people with diabetes, increasing the risk of medication errors and healthcare costs, and negatively impacting on individuals' ability to self-manage their condition (Flanagan et al, 2018).

Evidence demonstrates that optimising diabetes control helps to prevent long-term complications (UK Prospective Diabetes Study Group, 1998). These findings underscore the importance of embedding diabetes self-management education (DSME) within routine care to optimise long-term outcomes and reduce the burden on the NHS. Structured education programmes for people with type 2 diabetes are considered an integral part of diabetes care and should be delivered following diagnosis, meeting the individual's cultural, linguistic, cognitive and literacy needs (NICE, 2023).

There is a wealth of evidence validating the benefits of group-based DSME compared with

Citation: Whittle S, Bailey O, Partridge H, James J (2026) Type 2 diabetes rehabilitation: an exercise and education programme. *Journal of Diabetes Nursing* [Early view publication]

Article points

1. Combining structured exercise with diabetes self-management education improves HbA_{1c}, cardiovascular risk factors and mental wellbeing.
2. The 8-week Refocusise programme produced immediate reductions in HbA_{1c}, blood pressure, BMI and cholesterol.
3. Improvements in glycaemic control were sustained at 1-year and 2-year follow-up.
4. Integrating supervised exercise to diabetes education supports engagement, behaviour change and long-term self-management.

Key words

- Diabetes rehabilitation
- Education
- Exercise
- Self-management

Authors

Samantha Whittle, Bournemouth Diabetes Centre, University Hospitals Dorset NHS Foundation Trust; **Olivia Bailey**, Bournemouth Heart Club; **Helen Partridge**, Bournemouth Diabetes Centre, University Hospitals Dorset NHS Foundation Trust; and **Janet James**, Faculty of Health, Environmental and Medical Science, Bournemouth University.

“The management of chronic diseases such as cardiovascular disease and chronic obstructive pulmonary disease has shown improved outcomes following the implementation of rehabilitation programmes.”

routine care for people with type 2 diabetes (Steinsbekk et al, 2012). UK studies demonstrate that structured self-management programmes, including education and digital interventions, empower individuals to take an active role in managing their condition and result in modest but clinically meaningful improvements in HbA_{1c} (2.2–6.6 mmol/mol) at 6 months post-intervention. The most effective programmes are multi-component and involve enhanced contact time (>10 hours; Captieux et al, 2018).

There are several diabetes self-management programmes available in the UK, including DESMOND (Diabetes Education and Self-Management for Ongoing and Newly Diagnosed), the X-PERT Diabetes Programme and, locally in Dorset, the N2T2 (New to Type 2) education programme and REFOCUS (On Your Diabetes). Over time, the delivery of these programmes has diversified, with provision offered in both group and digital formats to accommodate varying patient needs and preferences.

Research indicates that these multi-component educational interventions can improve patient knowledge, psychosocial outcomes and health behaviours, although statistically significant improvements in HbA_{1c} are not universally reported (Deakin et al, 2006; Khunti et al, 2012; Chatterjee et al, 2018). Evidence from the DESMOND and X-PERT trials demonstrates improvements in illness beliefs, empowerment and lifestyle behaviours, with modest but clinically relevant reductions in HbA_{1c} in some cohorts (Deakin et al, 2006; Khunti et al, 2012; Wheatley et al, 2021). Locally developed programmes, such as N2T2, align with these principles, aiming to provide accessible education soon after diagnosis (Wheatley et al, 2021). However, while these programmes often discuss and promote increased physical activity, there is limited evidence of the effectiveness of those that include structured exercise as part of the programme.

The management of chronic diseases such as cardiovascular disease and chronic obstructive pulmonary disease has shown improved outcomes following the implementation of rehabilitation programmes (Richardson et al, 2019). Exercise and increasing physical activity, alongside improvements in lifestyle and diet, are key components of both cardiac and pulmonary

rehabilitation programmes. Physical activity, including planned and unplanned exercise, has been shown to greatly improve health outcomes and glycaemic management of individuals of all ages with type 2 diabetes (Kanaley et al, 2022).

However, people with type 2 diabetes are less likely to engage in physical activity than the general population (Morrato et al, 2007). A recent trial evaluating a structured exercise programme for people with type 2 diabetes demonstrated improvements in HbA_{1c}, although these did not reach statistical significance (Mukherji et al, 2022). Notably, this trial did not include dietary or other interventions that would typically form part of a rehabilitation programme.

Refocusise: a new service intervention

The Refocusise programme was developed to support people living with type 2 diabetes to improve the management of their condition while engaging in regular exercise. The programme is run weekly from the Bournemouth Heart Club (BHC) and is supported by one diabetes specialist nurse from University Hospitals Dorset and two exercise therapists from the BHC. To date, twelve courses have been completed, with group sizes ranging from 11–15 participants.

Anyone with type 2 diabetes who wants to participate can be referred by their GP or practice nurse. They are subsequently invited to enrol in the 8-week course that combines structured exercise sessions with an established DSME programme at the BHC. Baseline and post-intervention assessments are undertaken and include HbA_{1c}, blood pressure, body mass index, waist circumference, lipid profile and psychological well-being (using the Hospital Anxiety and Depression Scale [HADS]).

On completion of the course, participants are invited to join the BHC or are encouraged to continue with other forms of structured exercise. HbA_{1c} is reassessed at 1-year and 2-year follow-up to evaluate long-term glycaemic outcomes. As this is a service evaluation, NHS ethical approval was not required. Participants consented for their data to be collected to enable evaluation of the service.

The exercise component of the Refocusise course incorporates a carefully structured, progressive circuit designed to enhance cardiovascular health

Box 1. Refocusise: weekly structure of the combined exercise and education programme.

Week	Duration	Weekly plan	Weekly theme
1	3 hours	Pre-assessments: HAD scores, BP, weight and step tests; 1 hour diabetes education; 45 minutes exercise	Introduction to diabetes, signs and symptoms, effect on vascular health
2	2 hours	1 hour diabetes education 1 hour exercise	Eating to help manage blood glucose and health
3	2 hours	1 hour diabetes education 1 hour exercise	Diabetes treatments and medications
4	2 hours	1 hour diabetes education 1 hour exercise	Possible complications and annual reviews
5	2 hours	1 hour diabetes education 1 hour exercise	Eating to keep a healthy heart
6	2 hours	1 hour diabetes education 1 hour exercise	Theory of exercise and continuation strategies
7	2 hours	1 hour diabetes education 1 hour exercise	Digital technology/apps; health coaching opportunities A visit from LiveWell Dorset and the opportunity to sign up for health coaching/smoke stop and help with alcohol
8	3 hours	Post-course assessments: HADS/PAID scores and step tests	Final quiz, review and goal setting

“The circuit accommodates participants of varying abilities by offering both seated and standing exercise options, enabling individuals to engage at a level aligned with their comfort and capacity each week.”

while promoting overall functional strength in both the upper and lower body (*Box 1*). To ensure inclusivity, the circuit accommodates participants of varying abilities by offering both seated and standing exercise options, enabling individuals to engage at a level aligned with their comfort and capacity each week.

Specialist instructors deliver multiple intensity variations for each exercise, supporting gradual progression while preventing overexertion. Each movement includes weighted and body-weight alternatives, with clear demonstrations of how exercises can be modified to increase or decrease intensity as required.

Over the 8-week duration of the programme, the number of exercises performed is expanded to match participants' improving fitness levels. Higher-intensity options are introduced for those ready to progress, while individuals who begin with seated exercises are encouraged to transition toward standing positions where appropriate, fostering safe and sustainable advancement.

Results

The programme has been completed by 147 people (74 males), with a mean age of 64 years (range, 23–84 years). Initial results following the intervention have been positive, with improvements observed across all clinical markers (*Table 1*). These benefits were sustained over the longer term, with a mean reduction in HbA_{1c} of 11 mmol/mol at 1-year follow-up ($n=73$), which was maintained at 2 years following the intervention (*Table 2*).

Furthermore, the participants demonstrated improvements in mental wellbeing, as reflected in their HADS scores. Eighty-three participants completed the HADS at baseline and after programme completion. At baseline, the mean anxiety (HADS-A) score was 7.3 (SD, 4.2; range, 0–16), and the mean depression (HADS-D) score was 5.23 (SD, 3.1; range, 0–14). Post-programme, the mean anxiety score decreased to 5.8 (SD, 3.8; range, 0–16) and mean depression score to 3.6 (SD, 2.9; range, 0–10). HADS scores of 8–10 indicate mild

Table 1. Baseline and post-programme clinical markers.

Outcome (units)	Baseline (mean ± SD)	Post course (mean ± SD)	Mean difference (95% CI)
HbA _{1c} (mmol/mol) (n=147)	69 ± 16	60 ± 13	-9 (-11 to -7)
Total cholesterol (mmol/L)	4.3 ± 1.1	4.0 ± 1.0	-0.3 (-0.4 to -0.2)
BMI (kg/m ²)	33.2 ± 6.5	32.5 ± 6.3	-0.6 (-0.9 to -0.4)
Systolic blood pressure (mmHg)	139.0 ± 16.4	134.8 ± 14.7	-4.2 (-6.8 to -1.6)
Diastolic blood pressure (mmHg)	79.5 ± 8.8	75.7 ± 8.3	-3.8 (-5.5 to -2.0)
Oxygen saturation (%)	95.9 ± 1.9	97.0 ± 1.6	+1.1 (0.8 to 1.4)

BMI=body mass index; CI=confidence interval; SD=standard deviation fat.

Table 2. Longitudinal change in participant HbA_{1c} after the Refocusise programme.

Follow-up	HbA _{1c} (mmol/mol)	
	Mean	Mean change from baseline (95% CI)
1 year (n=73)	60	-11 (-14 to -8)
2 year (n=27)	59	-13 (-20 to -6)

CI=confidence interval.

anxiety or depression, 11–14 moderate and 15–21 severe symptoms (NICE, 2025).

Participants also provided written feedback describing the impact of the programme on their health and lives. One participant described Refocusise as “the start of the rest of your life”, while another reported that “This course has changed the way I think about diabetes.” Improvements in physical health associated with the exercise component were recognised through comments such as “My knees have improved” and “I can move my shoulder again”. Improvements in mental wellbeing were also noted, with comments such as “Everyone says I’m happier” and “I can play with the grandchildren more easily.”

Discussion

Evidence demonstrates that chronic disease management programmes (CDMPs) can positively impact the lives of people living with long-term conditions (Gbigbi-Jackson et al, 2024). Exercise-based cardiac rehabilitation is associated with reductions in cardiovascular mortality and hospitalisations, alongside improvements in quality of life (Dibben et al, 2023). Comparable benefits have been observed in exercise-based pulmonary rehabilitation, including improvements in respiratory symptoms and overall wellbeing (McCarthy et al, 2015).

Exercise is also a fundamental component of diabetes self-management. Both the American Diabetes Association (Colberg et al, 2016) and NICE (2026) recommend daily physical activity as a key lifestyle modification to support optimal diabetes control. Despite this, adults with type 2 diabetes often report higher levels of sedentary behaviour and lower engagement in physical activity compared to those without the condition (Kennerly and Kirk, 2018). Following completion of the Refocusise programme, a number of participants have joined the Bournemouth Heart Club and have continued to exercise on a regular basis, suggesting that participation may support longer-term engagement in physical activity.

The initial outcomes from the Refocusise programme indicate improvements across all measured clinical markers. Notably, the sustained reduction in HbA_{1c} suggests that this intervention may confer health benefits and, potentially, economic advantages. Evidence consistently shows that improved glycaemic control in people with diabetes is associated with significant cost savings, as even modest reductions in HbA_{1c} can reduce complications and subsequent healthcare expenditure (Baxter et al, 2016). Collectively, these findings reinforce the premise that early intervention and preventative strategies can be both clinically effective and economically beneficial.

The primary aim of Refocusise is to improve glycaemic control and promote holistic health and wellbeing in people with diabetes. Findings from this service evaluation suggest that these objectives were achieved within this small cohort. The next step is to extend the programme to a larger and more diverse population to determine whether these

outcomes can be replicated. This approach aligns with a Diabetes UK position statement emphasising the need for research into integrating structured exercise into diabetes care, drawing on established cardiac and pulmonary rehabilitation models (Morris et al, 2023).

Limitations

This service evaluation involved a small sample of participants who were referred by their GP and were aware that the programme included exercise, introducing a potential element of selection bias. Furthermore, the exercise programme was delivered by highly experienced staff at the Bournemouth Heart Club, a specialist facility that may not be accessible to the wider population should the intervention be scaled up. While the results are promising, they are derived from a relatively small and specific cohort. Future research should evaluate the effectiveness of this model in larger, more diverse cohorts and across different healthcare settings.

Conclusion

This service evaluation indicates that incorporating structured exercise into an enhanced diabetes self-management education programme can result in clinically meaningful improvements in glycaemic control, cardiovascular risk factors and psychological well-being. These benefits were observed immediately following programme completion and were sustained at one and two years. The findings support the integration and wider implementation of combined exercise and education interventions within routine diabetes care. ■

- Baxter M, Hudson R, Mahon J et al (2016) Estimating the impact of better management of glycaemic control in adults with Type 1 and Type 2 diabetes on the number of clinical complications and the associated financial benefit. *Diabet Med* **33**: 1575–81
- Captieux M, Pearce G, Parke HL et al (2018) Supported self-management for people with type 2 diabetes: A meta-review of quantitative systematic reviews. *BMJ Open* **8**: e024262
- Chatterjee S, Davies MJ, Stribling B et al (2018) Real-world evaluation of the DESMOND type 2 diabetes education and self-management programme. *Practical Diabetes* **35**: 19–23
- Colberg SR, Sigal RJ, Yardley JE et al (2016) Physical activity/exercise and diabetes: A position statement of the American Diabetes Association. *Diabetes Care* **39**: 2065–79
- Deakin T, Cade J, Williams R et al (2006) Structured patient education: The Diabetes X-PERT Programme makes a difference. *Diabet Med* **23**: 944–54
- Diabetes UK (2024) *How many people in the UK have diabetes*. DUK, London. Available at: <https://www.diabetes.org.uk/about-us/about-the-charity/our-strategy/statistics> (accessed 10.04.26)
- Dibben GO, Faulkner J, Oldridge N et al (2023) Exercise-based cardiac rehabilitation for coronary heart disease: a meta-analysis. *Eur Heart J* **44**: 452–69
- Dorset HealthCare University NHS Foundation Trust (2026) *Healthcare professionals*. Available at: <https://www.dorsethealthcare.nhs.uk/our-services-and-sites/physical-health/diabetes/healthcare-professionals> (accessed 08.02.26)
- Flanagan D, Dhataria K, Kilvert A (2018) Self-management of diabetes in hospital: A guideline from the Joint British Diabetes Societies (JBDS) for Inpatient Care group. *Diabet Med* **35**: 992–6
- Fowler MJ (2008) Microvascular and macrovascular complications of diabetes. *Clin Diabetes* **26**: 77–82
- Gbigbi-Jackson GA, Aguguo JC, Okobi OE et al (2024) Understanding the chronic disease management programs: a comprehensive literature review. *Medical Research Archives* **12**. Available at: <https://esmed.org/MRA/mra/article/view/5647> (accessed 26.11.25)
- Hex N, Bartlett C, Wright D et al (2012) Estimating the current and future costs of Type 1 and Type 2 diabetes in the UK, including direct health costs and indirect societal and productivity costs. *Diabet Med* **29**: 855–62
- Kanaley JA, Colberg SR, Corcoran MH et al (2022) Exercise/physical activity in individuals with type 2 diabetes: A consensus statement from the American College of Sports Medicine. *Med Sci Sports Exerc* **54**: 353–68
- Kennerly AM, Kirk A (2018) Physical activity and sedentary behaviour of adults with type 2 diabetes: a systematic review. *Practical Diabetes* **35**: 86–9
- Khunti K, Gray LJ, Skinner T et al (2012) Effectiveness of a diabetes education and self-management programme (DESMOND) for people with newly diagnosed type 2 diabetes mellitus: Three year follow-up of a cluster randomised controlled trial in primary care. *BMJ* **344**: e2333
- McCarthy B, Casey D, Devane D et al (2015) Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* **2**: CD003793
- Morrato EH, Hill JO, Wyatt HR et al (2007) Physical activity in U.S. adults with diabetes and at risk for developing diabetes, 2003. *Diabetes Care* **30**: 203–9
- Morris A, Bright C, Cocks M et al (2023) Recommendations from Diabetes UK's 2022 diabetes and physical activity workshop. *Diabet Med* **40**: e15169
- Mukherji AB, Lu D, Qin F et al (2022) Effectiveness of a community-based structured physical activity program for adults with type 2 diabetes: A randomized clinical trial. *JAMA Netw Open* **5**: e2247858
- NHS Digital (2016) *National Diabetes Audit, 2015-16 Report 2a: Complications and Mortality (complications of diabetes)*. Available at: <https://bit.ly/3QWP6ym> (accessed 10.04.26)
- NHS Digital (2020) *National Diabetes Inpatient Audit England and Wales (NaDIA) – 2019*. Available at: <https://bit.ly/4dglY0> (accessed 10.04.26)
- NICE (2023) *Type 2 diabetes in adults (QS209)*. NICE, London. Available at: <https://www.nice.org.uk/guidance/qs209> (accessed 10.04.26)
- NICE (2025) *Depression: Clinical Knowledge Summary*. Available at: <https://cks.nice.org.uk/topics/depression/> (accessed 08.02.26)
- NICE (2026) *Type 2 diabetes in adults: management (NG28)*. Last updated 29 June 2022. Available at: <https://www.nice.org.uk/guidance/ng28> (accessed: 26.11.25)
- Richardson CR, Franklin B, Moy ML, Jackson EA (2019) Advances in rehabilitation for chronic diseases: Improving health outcomes and function. *BMJ* **365**: l2191
- Steinsbekk A, Rygg L, Lisulo M et al (2012) Group-based diabetes self-management education compared to routine treatment for people with type 2 diabetes mellitus: A systematic review with meta-analysis. *BMC Health Serv Res* **12**: 213
- UK Prospective Diabetes Study (UKPDS) Group (1998) Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* **352**: 837–53
- Wheatley S, Arjomandkhan N, Murdoch C et al (2021) Improved blood glucose control, cardiovascular health and empowerment in people attending X-PERT structured diabetes education. *Practical Diabetes* **38**: 31–5
- Whicher CA, O'Neill S, Holt RIG (2020) Diabetes in the UK: 2019. *Diabet Med* **37**: 242–7

“Findings from this service evaluation suggest that the primary aims of improving glycaemic control and promoting holistic health and wellbeing in people with diabetes were achieved.”