
# Regional Diabetes Foot Pathway

Working together Excellence Openness & Honesty Compassion

**HSC Values** 

Jennifer Madden Consultant Podiatrist in Diabetes











### **The Diabetic Foot**

Triad of: Neuropathy Ischaemia

Infection










# Foot Assessments





# Neurological Assessment





Sensory assessment involves testing these sites

With 10 g Monofilament and Neurotip



# Vascular Assessment



### **PULSE PALPATION.**

DOPPLER ABPI ASSESSMENT SIGNS AND SYMPTOMS.





### Important to check both feet...



- PAD/CLI can occur unilaterally:
- Left foot palpable and Biphasic DP & PT
- Right foot Non palpable Monophasic PT and a Nonpalpable and non-audible DP. Severe rest pain.
- Patient required a DSA and the hallux went necrotic and is waiting amputation, necrosis on 2<sup>nd</sup> dorsal aspect PIPJ hopeful to save this toe





### <u>Risk tool</u>

#### Low Risk

No Risk Factors

Refer to podiatry for annual diabetes screening

#### Moderate Risk

One Risk Factor

Refer to podiatry for 6 monthly reviews

#### High Risk

Two or more Risk Factors

Refer to podiatry for regular review and routine care where needed usually 3-4 months

#### Active Risk

Ulcer, Charcot, Infection, Hot Red Swollen Foot

Urgently refer to podiatry for urgent assessment, active management and onward referral to MDFT by podiatry

# Proportion of population to foot risk

Total adult population with diabetes	Diabetic foot disease risk (NICE, 2015; SIGN, 2017)
/1-4%	Patients with active diabetic foot disease: presence of ulceration, or spreading infection, or critical limb ischaemia, or gangrene, or suspicion of an acute Charcot neuro-osteoarthropathy, or unexplained hot, red foot with or without pain.
4-8%	<b>High risk:</b> Patients with a history of diabetic foot ulceration or amputation, or more than one risk factor (e.g. loss of sensation, signs of peripheral arterial disease with callus or deformity, on renal replacement therapy).
20%	Moderate risk: Patients with one established risk factor for diabetic foot disease (e.g. loss of sensation, signs of peripheral arterial disease without callus, or deformity).
70%	<b>Low risk:</b> Patients at low risk of diabetic foot disease with no symptoms (apart from callus alone).

Working together

Excellence Openness & Honesty Compassion










# Low risk of foot ulcers and amputation








## Low risk of amputation and ulceration

- Advised on how diabetes may impact their feet
- Personal foot care does not need provided by podiatry. They can self-care for feet. Educate regarding their current foot risk
- Provided with contact of local podiatry foot protection team
- Enable to monitor feet or have carers monitor feet
- This is the majority of patient living with diabetes
- Made aware of what is urgent/emergency to contact podiatry or seek emergency advice over a weekend
- Currently biennial screening.







# Example of advice given

#### Tips for promoting healthy feet:


- Wash feet with soap and water and dry thoroughly
- ➔ Keep toenails trimmed and filed
- ➔ Apply moisturising cream daily
- ➔ Ensure the feet are clean
- Encourage good fitting shoes to be worn daily
- Avoid soaking the feet or applying moisturising cream between the toes as this can cause the skin to dry out or lead to cracks and cuts





#### Footwear should be:

- ➔ Worn daily
- → Well fitting
- Have a supportive heel counter and secure fastenings
- Checked regularly for a good fit, and any wear and tear
- Slippers should only be worn for short periods



If you notice any of the following foot health issues please speak to a healthcare professional e.g. Podiatrist, Medical Staff and/or Nursing Staff:

- → Skin breaks that are not healing
- ➔ Pus or discharge
- Any redness, discolouration, heat or swelling
- ➔ If a foot is colder and or paler than normal



Working together

Excellence

### Moderate Risk foot ulceration or amputation

✓ Signs of Neuropathy <u>or</u> PAD

To be seen 3-6 months by <u>Foot</u> <u>Protection Team</u>

Patient educated regarding risk to foot & foot health plan.

Record Risk

Pt can self refer to Podiatry Service







### High Risk

- ✓ History of ulceration or amputation
- ✓ PAD and neuropathy
- ✓ Callous or Deformity in presence of PAD and/or neuropathy



- Educate Patient regarding foot risk & foot health plan
- Record Risk
- Patient can self-refer to Podiatry service



# Active foot risk – all tiers of pathway

- Foot ulceration
- Infection
- Charcot neuroarthropathy
- Red hot swollen foot
- Wound care classification SINBAD
- Management of mild and moderate infection with oral antibiotics
- Offloading





# Why the regional diabetes foot pathway?





## Diabetes – Northern Ireland



All the people with diabetes in Northern Ireland could fill Windsor Park more than five times over There are over 108,000 people living with diabetes in Northern Ireland and this number is set to continue to rise.


Diabetes prevalence in Northern Ireland is growing around 4% year on year and with an estimated 376,000 people living here at risk of type 2 diabetes

(Diabetes UK 2022)





# Diabetic complications estimated costs



Major diabetes complications



COMPLICATION	COST
Stroke	£288 M
Kidney failure	£514 M
Ischaemic heart disease	£510 M
<b>Myocardial Infarction</b>	£603 M
Foot ulcers and amputations	<b>£986 M</b> Kerr,2012
Working together Excellence Openness & Hon	Compassion

### Foot complications in diabetes

In the United Kingdom, people with diabetes account for more than 40% of hospitalisations for major amputations and 73% of emergency room admissions for minor amputations.

We need to ensure that people access or are referred for treatment in and the right time, at the right place, and with the appropriate healthcare professional.





# Proportion of population to foot risk

Total adult population with diabetes	Diabetic foot disease risk (NICE, 2015; SIGN, 2017)
1-4%	Patients with active diabetic foot disease: presence of ulceration, or spreading infection, or critical limb ischaemia, or gangrene, or suspicion of an acute Charcot neuro-osteoarthropathy, or unexplained hot, red foot with or without pain.
4-8%	<b>High risk:</b> Patients with a history of diabetic foot ulceration or amputation, or more than one risk factor (e.g. loss of sensation, signs of peripheral arterial disease with callus or deformity, on renal replacement therapy).
20%	Moderate risk: Patients with one established risk factor for diabetic foot disease (e.g. loss of sensation, signs of peripheral arterial disease without callus, or deformity).
70%	<b>Low risk:</b> Patients at low risk of diabetic foot disease with no symptoms (apart from callus alone).





# Aim of Diabetic foot care pathway

- Better Outcomes for the Person Living with Diabetes
- Integrated and seamless
- Strengthen already established services
- Reflect regional strategic change
- Strong user voice
- Deliver NICE guidance





### Northern Ireland Diabetes Foot Care Pathway

#### The Diabetes Foot Care Pathway for Northern Ireland offers an improved service for adults living with diabetes.

#### Step 1: Foot Screening

- If you already live with diabetes or have been recently diagnosed, you will be assessed by a podiatrist/ podiatry assistant and an individual care plan will be agreed with you.
- You will be screened once a year to check your foot health.
- You will receive information on who to contact should you have any concerns about your feet.
- If problems are identified during your review you will be referred to step two of the pathway.



#### Step 2: Foot Protection Team (community based)

- This community based team provides early treatment and advice when a problem is identified.
- You will see a podiatrist at your local centre.
- A treatment plan will be discussed and agreed between you and your podiatrist.
- If your problem doesn't improve you will be referred to step three of the pathway.





#### Step 3: Enhanced Foot Protection Team (hospital based)

- Specialist hospital support is provided at this stage.
- You will see a podiatrist at your local hospital.
- Your problem will be continually assessed and treatment delivered by the most appropriate team.
- If you require further specialist treatment you will be referred to step four of the pathway.



#### Step 4: Multi-disciplinary Diabetes Foot Team (based at the Royal Victoria Hospital)

- Your local team will discuss your foot health with the regional Multi-disciplinary Diabetes Foot Team.
- You may be seen by the team in the Royal Victoria Hospital, Belfast.
- Foot problems that need treatment fast or are complex will be treated by this team.

If you have any concerns please contact your local podiatrist.

The Diabetes Network as an action from the Diabetes Strategic Framework for Northern Ireland launched a new Foot Care Pathway to improve services for adults living with diabetes in November 2019.

- The new Diabetes Foot Care Pathway enables all adults with diabetes to access the same services no matter where they live in Northern Ireland.
- The pathway encompasses routine screening, early treatment and care, specialist hospital support, and care when a person needs it from the Multi-Disciplinary Diabetes Foot Team based in the Royal Victoria Hospital in the Belfast HSC Trust.
- <u>Diabetes Foot Care Pathway HSCB (hscni.net)</u>





### <u>The Journey</u>

### November 2019

Launch of diabetes foot pathway

### March 2020 COVID19

Pathway continued to see patients with active foot risk

Including foot emergency/MDFT need

### May 2021


Rebuild starts – starting to tackling waiting list, restarting screening

### 2022

Continues rebuild and waiting list management MDFT rebuild 2023







HSC

### <u>Community foot</u> <u>protection role</u>

Biennial screenings for everyone with diabetes

# Offload at risk areas of the foot

ALL 5 trust within N Ireland have a community foot protection team

Referral via CGG accepted as well as self referral.

### Education

Onward referrals e.g. footwear, GP for CV or Diabetes, Refer further up pathway as needed

Podiatry treatment for high-risk patients Management of active risk patients




**HSC Values** 

HSC Health and Social Care

# **Biennial Foot Screening**









# Screening 2 year assurances requested from SPPG/DOH

Safety; amputation data and SAIs/HCAIs relevant to the change in screening arrangement. Experience; the experience of PLWD when accessing the new model of care. This work will be a coproduction with Diabetes UK, as part of a research project.





# Screening 2 year assurances requested from SPPG/DOH

Impact; the number of PLWD who are low risk and develop active foot disease within 12 or 24 months of their screening appointment.

HSC Health and Social Care Equity; the number of PLWD who when offered an appointment do not reply to a partial booking letter or who DNA their appointment.



### Moderate risk of foot ulceration or amputation


Attends foot protection team at a minimum every 6 months (always a discussion with patient)

Footwear advice and offload any pressure areas with insoles Aware of urgent/emergency advice





### High Risk of foot ulceration or amputation



Footwear discussed and offloading with insole if agreeable

Referral to footwear clinic as needed

Advice on emergency/urgent issues and how to contact







### EFPT

### Based or Linked to local N Ireland Hospitals.

Antrim Area, Altnagelvin Area, Causeway, Craigavon Area, Daisy Hill, South Western Acute and Ulster Hospital.



First contact for FPT and GP if urgent issue.

# Will assess non healing wounds

Can immediately refer to MDFT if emergency

Can admit and start IV antibiotics

For Belfast HSC FPT-MDFT fufils this element for patients



### Foot Ulceration

- Foot ulceration is a major complication of Diabetes
- Approximately 15% of all people with diabetes will be affected by a foot ulcer during their lifetime – 36.9m today and 57m by 2025 – conservative figures.
- Lifetime incidence rate of foot ulceration estimated between 19- 34%
- After healing recurrence rate of DFU is 40% within a year and 65% within 3 years
- 80% of amputations are preceded by foot ulceration
- Between 100-200 major/minor Amputations are carried out in NI every year
  Health and
  Social Care

### Regional Foot Protection Teams



# **Enhanced Foot Protection Teams**



MDFT

## Regional **Centre for** Endocrinology & Diabetes **Regional Diabetes Foot Centre**





Working together Excellence Openness & Honesty

# Early ulcer

# management






# Time is our Enemy





### MDFT, Level 1 regional diabetes foot unit. RVH








#### **Active Diabetic Foot Disease**

#### Diabetic Foot Emergencies Limb or Life Threatening





### EMERGENCY CRITERIA:

- Abscess,
- gas in tissue,
- wet gangrene,
- rapidly deteriorating CLI
- and/or systemic sepsis secondary to foot infection.
- Any individual with foot cellulitis that may need infection managed.




### **Urgent** Criteria

- Ulcer with PAD,
- Ulcer with suspected osteomyelitis,
- Suspected Charcot
- Also included in urgent referral are those patient with re-ulceration with deformity
- (Consolidated Charcot)
- stable dry gangrene
- and/or non-healing ulcer greater than 12 weeks if seeing EFPT (6 weeks Belfast FPT).
- These may initially be discussed at virtual MDFT/FPT huddle dependent on severity.





#### **Evolution of a Diabetic Foot Infection**



### Palliative/End of life care

Opinion can be sought regarding patient

Sending to Emergency department is to be avoided

Advice from Enhanced Foot protection or in most cases MDFT can be sought and recommendation placed on ECR/Encompass


Virtual clinics on complex frailty cases being piloted.







### Foot complications in diabetes



Five-year mortality and direct costs of care for people with diabetic foot complications are comparable to cancer

The mortality rate for people who undergo lower extremity amputation due to a DFU remains alarming:

More than half of people with a major amputation will be dead in 5 years . 5 year mortality for Charcot (29%), DFU (30.5%), minor (46.2%) and major amputations (56.6%) (Armstrong 2020)







Every 20 seconds a leg is lost to diabetes somewhere in the world.

Or in the time I have been speaking to you 90 major amputations!





### Take home message

Advocate	Highlight	Empower
Advocate for people living with diabetes to attend podiatry if and when invited to attend podiatry.	Always highlight what the urgent/emergency issues can be with their foot.	Empower people living with diabetes. The majority will be at low risk. Personal foot care is not required via a podiatrist.





## REFERRAL





# Belfast HSC Trust Foot protection team

BHSCT accept referrals from all health care professionals, family members, carers and the person themselves. Referral is made by telephone. Numbers listed below

South and East Belfast Area 02896158200

North and West Belfast area 02896158100





# Northern HSC Trust Foot protection team

NHSCT accept referrals from all health care professionals, family members, carers and the person themselves. The person making the referral should contact their nearest clinic. Their name and address is taken and a referral form will be posted to them for completion. Clinic contact numbers listed below

Larne Health Centre 028 28 261971 Carrickfergus Health Centre 028 93315811 Whiteabbey Health Centre 028 90808263 Glengormley Health Centre 028 90831410 Ballyclare Health Centre 028 93322575 Ballymena Health & Care Centre 028 25635672Thompson House, MUH Site 028 79300165 Cookstown Health Centre 028 86723811 Coleraine Community Services 028 70344831





# South Eastern HSC Trust Foot protection team

Self-referral is open to everyone. The process requires completion of a form that can be downloaded from the Trust's website or collected from the person's local clinic. When completed it is returned to the clinic or emailed to the central booking office. <u>CBO.podiatry@setrust.hscni.net</u>





# Southern HSC Trust Foot Protection team

SHSCT accept self-referral for new patients. Referral forms can be accessed by contacting CBU- Central booking Unit, 02837563444 and returned to CBU by email: <u>AHP.CBU@southerntrust.hscni.net</u>

Or post: AHP Central Booking Unit, Magowan West, 11,West Street, Portadown, BT62 3PG

New patient referrals can also be made by GPs, nursing and the allied health professions.





# Western HSC Trust Foot Protection team

All persons self-refer by completion of an application form. The form is emailed or posted to the person on request. Clinic contact numbers listed below

Woodview, Gransha - 02871865100

South West Area Hospital 02866382000

Omagh Hospital and Primary Care Complex 02882833100





#### Enhanced Foot protection Team – EFPT Monday to Friday 8.30 to 4.30 pm

Northern Antrim EFPT – 028 94424026 Causeway EFPT – 028 70346070 South Eastern Ulster - 07713008194 Southern **Central booking** Unit ,02837563444

#### Western

Woodview, Gransha- 02871865100 South West Area Hospital- 02866382000 Omagh -02882833100

**Belfast** MDFT covers EFPT component of pathway

# For Emergency 8.30 to 4.30 pm Monday to Friday

## **MDFT**

# 02896151444

Redirect to RVH Vascular specialist registrar out of hours (including bank holidays) and at weekends





