

A reliability study of wound assessment tools for diabetic patients in Indonesia

Suriadi Jais, Tisa Gusmiyah, Supriadi Syafei, Tuter Kardiatur, Hartono Maktal, Suhaimi Fauzan

Wound assessment tools specifically for diabetic foot wounds are less well developed than those for wounds such as pressure ulcers. The aim of this study was to evaluate the inter-rater reliability of a new wound assessment tool developed for diabetic foot wounds in Indonesia. The results showed that the inter-rater reliability of the MUNGS (maceration, undermining/tunnelling, necrotic tissue, granulation, signs or symptoms) tool was higher than that of a photographic wound assessment tool. The MUNGS tool may help monitor the progression of chronic wounds, especially among people with diabetes.

Indonesia is the world's fourth most populous country, with a population of 237.6 million in 2010 (Statistics Indonesia National Population and Family Planning Board Ministry of Health, 2012). According to a study of diabetes prevalence, Indonesia ranks seventh highest globally, with 7.6 million patients recorded in 2012, despite a relatively low prevalence of the disease (Soewondo et al, 2013). The two regions with the highest prevalence of diabetes in the country are Ternate, a small, remote island in eastern Indonesia, where 19.6% of the suburban population has diabetes, and the province of West Kalimantan, with a prevalence of 11.1% (Soewondo et al, 2013).

Diabetes is associated with multifaceted complications. One of the most common of these is foot ulcers, which often result in lower extremity amputations (Singh et al, 2005), and which have a prevalence of 4–10% among people with diabetes (Wu et al, 2007). Diabetic foot ulcers affect patients' quality of life and have social and economic consequences (Deribe et al, 2014). Diabetes-related complications increase the length of hospital stays and, therefore, the cost of care (Cichero et al, 2013). The average cost of an ulcer in a person with diabetes has been estimated at \$13,179 per episode (Stockl et al, 2004).

The high costs of treating diabetic ulcers emphasise the value of intensive inpatient or outpatient interventions designed to prevent ulcer progression. The management of wounds is an essential part of intensive care. In people with diabetes, inadequate treatment of diabetic foot ulcers will result in poor treatment outcomes (Itani et al, 2015). For example, the foot ulcer may become chronic and show delayed healing.

In recent years, interest in chronic wound management has increased, focusing not only on the correct treatment, but also on optimal preventive care (Restrepo-Medrano and Soriano, 2012).

Drawing on the evidence base, various wound management strategies have been developed to accelerate wound healing (Woodbury et al, 2004). The determinants of wound healing include the bioburden and severity of the wound, and these factors should be included in all wound assessments (Houghton et al, 2000).

To assess the effectiveness of wound treatment, a measurement tool is required that can describe the current condition of the wound and detect improvement or deterioration over time. The wound assessment process is essential for the development of an adequate treatment plan. Appropriate topical care and assessments of a wound are only possible when the observations and results of interventions are recorded.

Citation: Jais S, Gusmiyah T, Syafei S et al (2012) A reliability study of wound assessment tools for diabetic patients in Indonesia. *The Diabetic Foot Journal* 19: 95–99

Article points

1. The high costs of treating diabetic ulcers emphasise the value of intensive inpatient or outpatient interventions designed to prevent ulcer progression.
2. A cross-sectional pilot study of the inter-rater reliability of the MUNGS and PAWT tools was conducted at the Kitamura Wound Clinic in Pontianak city, Indonesia.
3. The inter-rater reliability of the MUNGS (maceration, undermining/tunnelling, necrotic tissue, granulation, signs or symptoms) tool was higher than that of a photographic wound assessment tool.

Key words

- Indonesia
- MUNGS tool
- Wound assessment

This article first appeared in *Wounds International* (2016) 7(2): 5–9

Authors

Suriadi Jais, Tisa Gusmiyah, Supriadi Syafei, Tuter Kardiatur, and Hartono Maktal, all study at the Institute of Nursing Muhammadiyah Pontianak, Kalimantan Barat, Indonesia; Suhaimi Fauzan studies at the School of Nursing, Faculty of Medicine, Tanjungpura University; Pontianak, West Kalimantan, Indonesia

Figure 1: The Assessment Tool for Diabetic Wound: MUNGS.

Patient's name: Date of admission:.....

	Score	Items	Score/Date
M		Maceration	
	0	None	
	1	Thin at the edge and/or maceration ≤ 2 cm from the wound edge	
	2	> 2 cm from the wound edge and/or expanded	
U		Undermining/tunnelling/sinus	
	0	None	
	1	≤ 3 cm	
	2	> 3 cm	
N		Necrotic tissue type (black, white, yellow, grey, brown, green)	
	0	None	
	1	Soft slough and with ≥ 1 colour	
	2	Necrotic; with spongy, soft and coloured skin	
	3	Necrotic; hard, spongy or moist tissue and skin with ≥ 1 colour	
	4	Necrotic; dry, hard, black and/or brownish	
G		Granulation tissue	
	0	Skin intact	
	1	Full granulation (100%)	
	2	Granulation of 50 % to $< 100\%$	
	3	Granulation of $< 50\%$	
	4	No granulation	
S		Other wound-related signs or symptoms	
	0	None	Wound edge: Around the skin wound:
	1	One or two	<input type="checkbox"/> Red ring <input type="checkbox"/> Hyperpigmentation
	2	Three to five	<input type="checkbox"/> Hyperkeratonic <input type="checkbox"/> Induration
	3	More than five	<input type="checkbox"/> Unattached <input type="checkbox"/> Hypopigmentation
			<input type="checkbox"/> Undefined <input type="checkbox"/> Erythema
			<input type="checkbox"/> Crust <input type="checkbox"/> Oedema
			<input type="checkbox"/> Pale <input type="checkbox"/> Purple
			<input type="checkbox"/> Damage <input type="checkbox"/> Lesion
			<input type="checkbox"/> Epibole Granulation:
			<input type="checkbox"/> Rolled/lining <input type="checkbox"/> Fragile granulation
			Wound infection or inflammation: <input type="checkbox"/> Bright red
			<input type="checkbox"/> Pain <input type="checkbox"/> Hypergranulation
			<input type="checkbox"/> Pus <input type="checkbox"/> Senescent
			<input type="checkbox"/> Odour <input type="checkbox"/> Pale
<input type="checkbox"/> Fever <input type="checkbox"/> Blackish			
<input type="checkbox"/> Rising temperature/warm <input type="checkbox"/> Trauma			

Monitoring tools used in Indonesia include the leg ulcer measurement tool (LUMT) and the photographic wound assessment tool (PAWT) (Houghton et al, 2000; Woodbury et al, 2004). It is very important to observe diabetic ulcers at every change of dressing to determine ulcer progression and to identify potential barriers to wound healing and serious complications.

Developing a new tool

The authors have created a new assessment tool for evaluating wound healing progression. The MUNGS tool (**m**aceration, **u**ndermining/tunnelling, **n**ecrotic, **g**ranulation and **s**igns or symptoms) was developed based on the authors' clinical observations of diabetic ulcer patients in Indonesia (Figure 1).

The total MUNGS score for each wound is calculated by summing the scores assigned to each of the five domains. Thus, the range of possible total MUNGS scores is between 0 and 15, with 0 representing a completely healed ulcer, and higher scores indicating poor wound healing progress.

PAWT is the tool used in the authors' clinic currently. This is a modified version of the pressure sore status tool. The PAWT consists of six parameters: wound edges, necrotic tissue type and amount, skin colour surrounding wound, granulation tissue type, and epithelialisation.

There are a number of classification systems available to grade ulcers according to the presence and extent of various physical characteristics of the wound (International Best Practice Guidelines, 2013). However, the authors felt there was a need to develop a specific assessment tool for the purposes of ongoing monitoring of progression.

MUNGS is the first instrument developed in Indonesia specifically to evaluate the appearance of diabetic foot wounds. Unlike PAWT, MUNGS does not include an assessment of the wound size and depth. The rationale for this lies in the difficulties in gaining accurate measurement of the depth and size of diabetic foot ulcers, which can often be on the toes, where it is difficult to accurately measure the size of the wound. The MUNGS tool includes signs or symptoms and uses this as a parameter of healing progress rather than size.

Assessing the wound for clinical signs and symptoms of inflammation and infection is of particular importance in people with diabetes^[13], and

clinicians can use assessment tools such as NERDS (non-healing wounds; exudating wounds; red and bleeding granulation tissue; debris on wound surface (yellow/black); and smell) and STONES (size – bigger; temperature – increased; Os – probe to or exposed bone; new or satellite areas of breakdown; exudate, oedema, erythema; and smell). NERDS is for detection of superficial infection, while STONES is to detect deep infection.

The authors believe the parameters identified in the MUNGS tool include components that will recognise infection and ischaemia, but this will require further research to gain confidence in this aspect of assessment.

Prior to using an assessment tool, the reliability of the instrument must be evaluated. The goal of the present study was to compare the inter-rater reliability of MUNGS and PAWT in assessing wound healing progression in diabetic patients. Inter-rater reliability is the degree of concordance or consistency of the performance of two or more observers in recording the same responses at the same time (Karanicolas et al, 2009).

Method

A cross-sectional pilot study of the inter-rater reliability of the MUNGS and PAWT tools was conducted at the Kitamura Wound Clinic in Pontianak city, Indonesia.

Fourteen adult outpatients with diabetic ulcers were asked to participate in the evaluation of the assessment tools. The inclusion criteria were patients who were physically able to participate and whose diabetic ulcers included a variety of wound sizes. A pre-study sample size calculation indicated, that with 14 subjects, a two-tailed test, kappa ($P \geq 0.00$) and 80% power were detected (Sim and Wright, 2005).

Raters

Eight raters conducted the assessment of the diabetic wounds: four experienced wound care nurses (WCNs) with a minimum of 2 years' experience and four inexperienced raters. The inexperienced raters were final year undergraduate nursing students. The WCNs had previous training in diabetic wound care and did not receive any further instructions on how to use the assessment tools. The student nurses were educated about diabetic ulcers and how to use the assessment tools (MUNGS and PAWT) over 2 days. On the first day, students were taught the characteristics of the wound bed in diabetic foot ulcer using photographs,

References

- Bates-Jensen BM, Vredevoe DL, Brecht ML (1992) Validity and reliability of the Pressure Sore Status Tool. *Decubitus* 5: 20–8
- Bates-Jensen BM (1997) The Pressure Sore Status Tool a few thousand assessments later. *Adv Wound Care* 10: 65–73
- Cichero MJ, Bower VM, Walsh TP, Yates BJ (2013) Reducing length of stay for acute diabetic foot episodes: employing an extended scope of practice podiatric high-risk foot coordinator in an acute foundation trust hospital. *J Foot Ankle Res* 6: 47
- Deribe B, Kifle Woldemichael K, Namera G (2014) Prevalence and factors influencing diabetic foot ulcer among diabetic patients attending Arbaminch Hospital, South Ethiopia. *J Diabetes Metab* 5: 1
- Houghton PE, Kincaid CB, Campbell KE et al (2000) Photographic assessment of the appearance of chronic pressure and leg ulcers. *Ostomy Wound Manage* 46: 20–30
- International Best Practice Guidelines (2013) *Wound Management in Diabetic Foot Ulcers*. London: Wounds International
- Itani H, Gandoura NA, Ahmed T, Ahmed RM (2015) Impact of psychological stress on wound healing for patients with diabetic foot ulcers. *The Diabetic Foot Journal Middle East* 1: 18–22
- Karanicolas PJ, Bhandari M, Kreder H, et al (2009) Evaluating agreement: conducting a reliability study. *J Bone Joint Surg Am* 91(Suppl 3): 99–106
- Landis JR, Koch GG (1977) The measurement of observer agreement for categorical data. *Biometrics* 33: 159–74
- Pillen H, Miller M, Thomas J et al (2009) Assessment of wound healing: validity, reliability and sensitivity of available instruments. *Wound Practice and Research* 17: 208–17
- Restrepo-Medrano JC, Soriano JV (2012) Development of a wound-healing index for chronic wounds. *EWMA Journal* 12: 39–46

Table 1. Inter-rater reliability results for MUNGS.

Item	Weighted kappa		Confidence interval (95%)	
	Nurse	Student	Nurse	Student
All MUNGS items	0.81	0.69	0.71–0.92	0.52–0.86
Individual MUNGS items				
Maceration	0.93	0.14	0.82–1.0	–0.19–0.48
Undermining/tunnelling	1.0	1.0	1.0–1.0	1.0–1.0
Necrotic tissue type	0.88	0.47	0.71–1.0	0.16–0.77
Granulation tissue	0.88	0.42	0.73–1.0	0.10–0.71
Additional signs/symptoms	0.51	0.42	0.15–0.87	0.90–0.76

Table 2. Inter-rater reliability results for photographic wound assessment tool (PAWT).

Item	Weighted kappa		Confidence interval (95%)	
	Nurse	Student	Nurse	Student
All PAWT items	0.60	0.43	0.44–0.76	0.16–0.69
Individual PAWT items				
Edges	0.19	–0.14	–0.07–0.44	–0.34–0.06
Type of necrotic tissue	0.47	0.00	0.23–0.72	–0.20–0.20
Amount of necrotic tissue	0.54	0.25	0.33–0.75	–0.10–0.60
Skin colour surrounding the wound	0.40	0.05	0.10–0.71	–0.22–0.32
Granulation tissue	0.53	0.52	0.21–0.85	0.13–0.91
Epithelialisation	0.36	0.49	0.06–0.66	0.12–0.86

and on the second day they used the tools for assessment on three diabetic foot ulcer patients.

Study protocol

All patients in this study were present at the outpatient clinic during the data collection period and were recruited by the wound care nurse in the routine care process. The patients were informed about the study goals and procedures, and when they agreed to participate in the study they signed consent forms.

Two WCNs and two students rated the diabetic wounds using MUNGS, and two nurses and two students rated the wound using PAWT. The patients were assessed by the raters, and other wound care nurses who did not participate in this study changed the dressings. Each assessment was performed simultaneously and independently. Participants were blind to the ratings of the other evaluators. The total MUNGS and PAWT scores were calculated.

In all the assessments, precautions were taken to avoid cross-contamination of the wounds. Fresh

gloves were used during each evaluation, a waterless hand cleanser was used between the evaluations, and no measurement instruments were transferred between patients.

Statistical analysis

The inter-rater reliability was expressed in terms of Cohen's kappa coefficient. This is a measure of association that indicates the agreement of scores measured by two raters. For example, a value of 0.60 denotes acceptable agreement between assessors, and a value of 0.80 denotes satisfactory or good agreement.

The scale is: score of <0=poor, 0–0.20=slight, 0.21–0.40=fair, 0.41–0.60=moderate, 0.61–0.80=substantial and 0.81–1.00=almost perfect agreement (Landis and Koch, 1977). Cohen's kappa was calculated using MedCalc® version 15.8.

Results

In this study, wounds were assessed in 14 people with diabetes. The mean age of the patients was 54.6 years (SD 7.98; range 38–65 years), and 64.3% were women. The mean length of the wounds on the time of the assessment was 27.6 (SD 28.5; range 1–90 days). Each patient had one ulcer.

The inter-rater reliability of MUNGS among the WCNs was almost perfect agreement (0.81), whereas it was substantial (0.69) among the students (Table 1). For the individual MUNGS items used by the WCNs, agreement was satisfactory, except for signs or symptoms, where it was moderate (0.51); and for the students, each item was satisfactory, except for maceration, where it was poor (Table 1).

Using the PAWT, overall agreement was moderate for the WCNs (0.60) and fair for the students (0.43) (Table 2). There was moderate agreement for three of the PAWT items assessed by the WCNs, and for wound edges and epithelialisation the level of agreement was less than 0.40 (Table 2). Agreement for the PAWT items assessed by the students was moderate for granulation tissue (0.52) and epithelialisation (0.49). The level of agreement was below 0.40 for the necrotic tissue type (0.00), amount of necrotic tissue (0.25), skin colour surrounding the wound (0.05) and wound edges (–0.14). Wound edge had a negative rating — a negative kappa represents agreement worse than expected, or disagreement among raters. The kappa value of 0 for necrotic tissue type indicates no more rater agreement than that expected by chance.

Discussion

The results of this study demonstrated that the inter-rater reliability of the MUNGS tool was higher than that of the PAWT among both groups of raters (WCNs and students). Interestingly, the overall agreement of the PAWT was moderate for the two groups of raters, whereas the coefficients of agreement for most of the PAWT items were less than 0.60. This is probably due to the variation in the number of years of experience of the raters, especially that of the nurses.

Another study reported that the reliability of the PAWT was greater in individuals who had at least 5 years of clinical experience in chronic wound management (Houghton et al, 2000).

Based on the findings of this study, the authors can conclude that clinical experience provides the necessary training to distinguish various descriptors of a wound, such as the condition of the wound bed, the skin surrounding the wound and other signs or symptoms specific to diabetic wounds.

The MUNGS results showed that signs and symptoms was moderate for WCNs. This may be caused by several factors. In this study, the ability of two raters to obtain the same results when performing a clinical test can be affected by a number of signs and symptoms on wound condition and/or examiner error. For the item, examiner error could occur if signs and symptoms were identified and/or interpreted inconsistently, and the exam results may be different. The item maceration was poor for students. A possible explanation for this finding is years of clinical experience. The training period that was completed prior to data collection may have improved test performance. However, this training period needs to be considered when applying the results of this study in practice.

The need for prior clinical experience has also been reported with other wound assessment tools, such as the Pressure Sore Status tool (Bates-Jensen, 1992; 1997).

The coefficient of agreement for most of the MUNGS items was greater than 0.80 for the experts. The only mean coefficient of agreement below 0.80 for the WCNs was for additional signs or symptoms. Cohen's kappa value for inter-rater reliability was almost perfect (Landis and Koch, 1977).

Cohen's kappa value for the agreement among the students on the undermining/tunnelling MUNGS

items was perfect. This may be explained by the amount of undermining/tunnelling being easily identified and recorded in the assessment of wounds.

A previous study by Houghton et al of the use of PAWT for assessing leg ulcers reported that the inter-rater reliability of experienced individuals was high (Houghton et al, 2000). However, that study not only used photographs of diabetic ulcers, but also ischaemic and venous ulcers. Houghton et al postulated that the coefficients of inter-rater reliability were higher when the photographic assessment tool was applied to pressure ulcers compared to leg ulcers (Houghton et al, 2000).

The current study was based on diabetic ulcers of patients in a clinical setting and differs from Houghton et al in the clinical setting, type of wounds assessed and statistical analysis (Houghton et al, 2000). In practical situations many more elements can be taken into account, such as whether or not undermining/tunnelling and signs or symptoms in the wound are present. Wounds in people with diabetes may differ greatly between individuals.

Prior to their application in practice, all assessment tools must be shown to provide a reliable measure of wound healing (Pillen et al, 2009). As with other tools that have been developed to assess wound appearance, the MUNGS tool must be used by individuals with clinical experience. The reliability of this tool will be compromised in the hands of inexperienced trainees. Using the MUNGS to assess wound appearance has advantages, but has limitations as well. Responsiveness of the MUNGS has not been demonstrated by its ability to adequately detect change in diabetic ulcer appearance over time and to demonstrate a difference between healers and non-healers. In addition, aspects related to ischaemia need further development.

Conclusion

This cross-sectional pilot study showed that the inter-rater reliability of the MUNGS tool for assessing diabetic wounds was almost perfect. These findings provide evidence supporting the application of the MUNGS tool in the assessment of the progression and healing of wounds, especially diabetic wounds, of inpatients and outpatients. Further studies with larger groups of subjects are required to determine the validity of the study and the utility of the MUNGS tool over time. ■

- Sibbald RG, Woo K, Ayello EA (2006) Increased bacterial burden and infection: the story of NERDS and STONES. *Adv Skin Wound Care* **19**: 447-61
- Sim J, Wright CC (2005) The kappa statistic in reliability studies: use, interpretation, and sample size requirements. *Phys Ther* **85**: 257-68
- Singh N, Armstrong DG, Lipsky BA (2005) Preventing foot ulcers in patients with diabetes. *JAMA* **293**: 217-28
- Soewondo P, Ferrario A, Tahapary DL (2013) Challenges in diabetes management in Indonesia: a literature review. *Global Health* **9**: 63
- Stockl K, Vanderplas A, Tafesse E, Chang E (2004) Costs of lower-extremity ulcers among patients with diabetes. *Diabetes Care* **27**: 2129-34
- Statistics Indonesia National Population and Family Planning Board Ministry of Health (2013) *Indonesian Demographic and Health Survey 2012*. MEASURE DHS ICF International, Jakarta
- Woodbury MG, Houghton PE, Campbell KE, Keast DH (2004) Development, validity, reliability, and responsiveness of a new leg ulcer measurement tool. *Adv Skin Wound Care* **17**(4 Pt 1): 187-96
- Wu SC, Driver VR, Wrobel JS, Armstrong DG (2007) Foot ulcers in the diabetic patient, prevention and treatment. *Vasc Health Risk Manag* **3**: 65-76