

# Meeting NICE Gold Standards (2015): a feasibility study to improve podiatry accessibility for people with diabetes and foot ulcers

Timothy Simon Caslake, Charlotte Moen

**Citation:** Caslake TS, Moen C (2016) Meeting NICE Gold Standards (2015): a feasibility study to improve podiatry accessibility for diabetic patients with foot ulcers. *The Diabetic Foot Journal* 19: 26–30

## Article points

1. The key challenge discussed within this article is how to provide a better service for diabetic patients with foot ulcers. The solution was the introduction of a 'drop-in' clinic that aimed to provide patients instant access, meet the NICE minimum requirements of seeing the patients within 1 working day and is both clinically and cost effective.
2. The long-term impact of the 'drop-in' clinic was twofold; the quality of life for those patients with ulcers improved and there are potential savings of over £100,000 in amputation costs per hospital (Nason et al, 2012).
3. The service improvement project is an example of clinical leadership, the project being led by a band 7 podiatrist.

## Key words

- Cost savings
- Drop-in diabetic foot ulcer clinic
- Twenty-four hour access

## Authors

Timothy Simon Caslake, Podiatry team leader band 7, Southport and Ormskirk NHS Trust; Charlotte Moen, Programme Lead/Senior Lecturer Professional Development, Edge Hill University

**A person with diabetes and a skin lesion should be referred within one working day and triaged one working day later; this is the gold standard for NHS Trusts (NICE, 2015). The aim of this project was to improve the accessibility to the podiatry service for all patients with diabetes with ulcers and to meet the NICE standards (2015) for treatment of people with diabetes and skin lesions/ulcers. The objectives of the project were to pilot a 'drop-in' diabetic foot ulcer clinic, to audit the number and type of patients seen within the clinic, as well as the number of foot ulcer admissions to accident and emergency, and assess the impact of the 'drop-in' clinic on A&E. The impact of the service on those attending the clinic was also evaluated.**

Five per cent of people with diabetes in the UK experience a foot ulcer (NICE, 2015). Furthermore, "sixty per cent of diabetics, with foot ulcers develop infection and, as a result, up to twenty five per cent of these may have to undergo partial or full, lower-foot or lower-limb amputation," according to Mooney (2013). It is estimated that £650 million is spent on diabetic foot ulcers and amputations each year in the NHS (Kerr, 2012; NICE, 2015). However, it is unclear why 25% of these patients had an amputation; it could be argued this is due to infection (Turner, 2013) and/or because of a lack of rapid access pathways, as Nice Guideline 19 advocates referral to the multidisciplinary foot team (MDFT) within 24 hours of admission. Turner (2013) highlighted that 80% of amputations are preventable, with rapid access being one of the main contributors to prevention.

In the UK alone, 100 people with diabetes have a lower-limb amputation a week, with costs to the NHS amounting to as much as £700 million per year (Holman et al, 2012). The cost to treat a foot ulcer is estimated at between £3,000 and £5,000 (Fard et al, 2007), with diabetic foot-related problems accounting for more hospital admissions than all of the other diabetic-related problems combined (Yarwood-Ross and Randall, 2013). This is also supported by Kerr

(2012), who reported that the financial cost of diabetic foot care is between £1.75 million and £1.82 million per day.

Diabetes UK (2011), Moulton (2013) and NICE (2015) conclude that having a rapid access service can benefit the patient by reducing the risk of an ulcer developing further complications, such as infection, amputation and necrosis.

In response, diabetic foot screening has increased in trusts over the past year and this has been supported by a national initiative, the NICE guideline 'Put Feet First' (Turner, 2013).

## Challenges for the NHS

The challenge for the NHS is to make financial savings at the same time as trying to meet the NICE guidelines. Mahaffey et al (2012) found that people with diabetes are admitted to hospital twice as often as people without diabetes and they occupy beds twice as long. They also reported that a saving of £35,000 was made over a 3-year period when people with diabetes were referred to the diabetic specialist nurse in the community and there was also a reduction in the amount of patients being admitted to hospital.

## Current practice

Despite the high risk factors associated with diabetes

### Page points

1. This service improvement project demonstrates that it is possible to improve the quality of patient care and patient experience, while reducing costs.
2. Clinicians are in an ideal position to both lead and implement service improvement projects.
3. The project provides an example of an multidisciplinary team approach to service improvement based on collective leadership.

and ulcers, and the emphasis placed on the need for preventative measures, Trusts were found to not be adhering to the previous NICE 2011 guidelines (McInnes, 2012). In addition, Trusts are not doing enough to prevent amputation (Jeffcoate and Rayman, 2011; McInnes, 2012). Staffing issues, capacity and cost-saving initiatives are reasons that could be attributed for the lack of adherence.

Where Trusts have introduced clear pathways for people with diabetes with an active ulcer to have rapid access to the MDFT, a reduction of 50% of amputations in hospitals has been seen (Moulton, 2013). Moulton's (2013) findings are supported by North Mersey Diabetes Network (NMDN, 2011) and by the Diabetes UK (2013) report 'Fast Track for a Foot Attack: Reducing Amputations'.

### Aims

The aims of this pilot were to improve the accessibility to the podiatry service for all patients with diabetes with foot ulcers, and to meet the NICE NG19 standards (2015) for treatment of people with diabetes with skin lesions/ulcers.

### Objectives

- To pilot a 'drop-in' diabetic foot ulcer clinic
- To audit the number and type of patients seen within the 'drop-in' diabetic foot ulcer clinic
- To audit the number of foot ulcer admissions to A&E and assess the impact of the 'drop-in' clinic on A&E.

### Method

A pilot 'drop-in' clinic was opened for diabetic patients over an 8-week period (between December 2, 2014 and January 28, 2015). Eight emergency 'drop-in' appointments were made available between 8.30am and 11.30am on a Monday morning each week.

Data were collected from both the podiatry and A&E computer system for 8 weeks prior to the 'drop-in' clinic commencing and 8 weeks during the clinics. The number of foot ulcer admissions to A&E and to the podiatry department was recorded during this time period. In addition, a patient satisfaction survey was conducted at the end of each treatment during the pilot phase.

### Results

Forty patients were treated during the 8-week 'drop-

in' clinics. Seven had ulcers treated, one was pre-ulcerative, two had ingrowing toenails with infection, three had painful corns with risk of breakdown and the remaining 27 were deemed non-urgent.

Table 1 shows the patient satisfaction questionnaire that was handed out to patients, of which 80% (32/40) responded. It was also noted that all the respondents would recommend the service to family and friends.

## Discussion

### Benefits

Prior to the pilot, patients with ulcers firstly attended their GP surgery before being referred to the podiatry department. This resulted in a significant delay between the problem being detected and being treated. NICE NG19 standards (2015) and NICE Prevention and Management (2015) state that people with diabetes with a break in skin integrity must be seen by a specialist within one working day and this was achieved during the pilot. During this study, eight patients treated with ulcers were potentially prevented from developing further complications, such as amputation. Furthermore, by using the NICE NG19 standards (2015) as a template, the authors were able to produce a clear pathway and a standard operating procedure for diabetic foot ulcers.

Admissions of diabetic foot ulcers to A&E reduced during the pilot and, as a result, there were reduced waiting times and improved patient satisfaction within A&E. However, there were variables that could have affected the reliability of the data collected from A&E due to different computer systems operating within podiatry and A&E. This was a limitation of the project. Similar issues were also identified within the literature, for example, coding issues (Royal College of Nursing, 2013) and incompatible systems (Pope et al, 2013). This results in non-comparable data, which means impact data are difficult to extract. Therefore, the impact of the 'drop-in' clinic could be more significant than the figures suggest (i.e. the number of people with diabetes and ulcers could be under-recorded).

Patient satisfaction with the 'drop-in' clinic was high (91% rated care as 'excellent' and 9% as 'good'). An example of the clinical impact was a patient who, having discharged himself from hospital due to the treatment he received, attended the clinic with a seriously infected ulcer that could be probed to the bone. With treatment that included intravenous

antibiotics, a potential below-knee amputation was avoided. The action taken not only impacted positively on the patient's quality of life, but impacted positively in financial terms, as this represented a potential saving to the NHS of £35,000 on amputation costs (Mahaffey et al, 2012; Moulton, 2013). In total, eight ulcers were treated, which had they been left longer than 24 hours, might have deteriorated and developed further complications.

The long-term impact of the 'drop-in' clinic was twofold; the quality of life for those patients with ulcers improved and there are potential savings of over £100,000 in amputation costs per hospital (Nason et al, 2012). It was also noted that further savings could be made as people with diabetes are admitted to hospital twice as often as patients without diabetes (Mahaffey et al, 2012); they have extended stays in hospital with unknown associated costs (Stang and Munro, 2015) and it has been estimated that the cost to treat an individual ulcer is between £3,000–£5,000 (Fard et al, 2007). Furthermore, North Mersey Diabetes Network (NMDN, 2011) stated that when a patient is seen by a member of the multidisciplinary footcare services within the recommended 24-hour period, the chance of preventing further complications is increased. Therefore, the right care and treatment can reduce the risk of amputations and the length of hospital stay (NICE, 2015).

## Challenges

The unintended outcomes of the pilot were that 27 non-urgent patients were seen (representing 67% of the total number of patients). The reasons for this could be a lack of patient understanding about what is classed as 'urgent' for people diabetes and also the purpose of the 'drop-in' clinic (Diabetes UK, 2011; NICE, 2011; NMDN, 2011). In order to further develop the clinic and prevent this scenario from reoccurring, better patient education for people with diabetes is required, as well as clear inclusion and exclusion criteria for the 'drop-in' clinics.

A patient satisfaction survey reported that 87% of the patients expressed a need for the clinic more than once a week because of the rapid access. Patients reported themes such as: it was 'easier and better for them'; that they could just 'turn up' for an appointment; and that they found the current system of booking a routine treatment 'inefficient'. These findings resonate with patient complaints; the fact

**Table 1. Drop-in clinic' patient satisfaction questionnaire.**

Question	Yes	No
Do you think the 'drop-in' clinic should be more than once throughout the week?	87%	13%
Have you ever attended A&E for an ulcer on your foot?	16%	84%
Would you recommend this service to friends and family if they needed similar care or treatment?	100%	0%
Did you prefer attending a 'drop-in' clinic rather than making a set appointment?	56%	44%

**Table 2. Cost-effectiveness analysis of a 'drop-in' diabetic foot ulcer clinic.**

Variable	Description
Population	People with diabetes
Benefit	<ul style="list-style-type: none"> <li>■ Diabetic urgent referrals seen within the NICE gold standards (2011) 24 hours guideline</li> <li>■ Detection of early signs of deterioration (Diabetes UK, 2009; NICE, 2014 )</li> <li>■ Improved patient satisfaction</li> <li>■ Reduced cost of further stays in hospital</li> <li>■ Reducing risk of amputation (Diabetes UK, 2011; NICE, 2015)</li> </ul>
Harms	<ul style="list-style-type: none"> <li>■ No multidisciplinary foot team</li> </ul>
Net benefits	<ul style="list-style-type: none"> <li>■ Eight diabetic patients with foot ulcers treated</li> <li>■ Potential saving of £35,000 per amputation (Mahaffey, et al 2012; Moulton, 2013)</li> <li>■ Cost to treat an ulcer is between £3,000–£5,000 (Fard et al, 2007)</li> </ul>
Approximate costs	<ul style="list-style-type: none"> <li>■ Band 6 cost for 3 hours 30 minutes — £52</li> <li>■ Equipment and dressing per ulcer — £10</li> </ul>
Balancing net benefits with costs	<ul style="list-style-type: none"> <li>■ Cost to run clinic for 8 weeks — £600–700</li> <li>■ Potential benefits of treating 8 patients (x 8 amputations prevented £280,000 x8 cost to treat an ulcer between £24,000 and £40,000) — Total saving = £304,000</li> </ul>

that they 'find it hard to get an appointment on the same day they phone' (Murdock et al, 2014). It is worth noting that these reasons could be the catalysts that make such a high number of non-urgent patients attend the 'drop-in' clinic. It is clear that, in addition to further expanding the clinic, the booking system of routine treatment needs to be improved.

## Conclusion

The aim of improving the accessibility of the podiatry service for all people with diabetes and foot ulcers in order to meet NICE NG19 standards (2015) was achieved. This was evidenced through the audits completed prior and during the 'drop-in' clinics.

Patient satisfaction indicated positive experiences for all patients in terms of quality of care and treatment. There was a positive impact on the A&E department as there were fewer patients to treat and, thus, waiting times reduced. A clear pathway and standard operating procedure was established for diabetic foot ulcer patients. Furthermore, there are clear potential cost savings to be made, in terms of ulcer prevention/management and costly amputation, especially if clinicians have access to a multidisciplinary foot care service (NICE, 2015). ■

### Recommendations

Sixty-seven per cent of patients assessed within the 'drop-in' clinic were non-urgent. Therefore:

- A 'drop-in' clinic should be available for all existing diabetic patients who are concerned about their feet
- Structured education should be available to help patients understand their condition and the appropriate pathway for treatment
- It is recommended that a patient information leaflet is developed to explain the 'drop-in' clinic inclusion/exclusion criteria and what is classed as 'urgent'
- To extend the clinic to other long term conditions as recommended by the Department of Health (2008)
- Research is required to explore whether organisations are meeting the NICE NG19 standards (2015) for the treatment of people with diabetes who have a break in the skin and then enable the sharing of best practice.

Care Quality Commission, (2010) *The State of the Health Care and Adult Social Care in England: Key Themes and Quality of Services in 2009*. Available at: <http://bit.ly/1Riiet8> (accessed 11.01.2016)

Department of Health (2008) High Quality care for all NHS- the next stage review final report. Available at: <http://bit.ly/1gr7XEk> (accessed 11.01.2016)

Diabetes UK (2009) *Putting Feet First: Commissioning Specialist Services for the Management and Prevention of Diabetic Foot Disease in Hospitals*. Diabetes UK, London

Diabetes UK (2011) *Putting Feet First: Commissioning Specialist Services for the Management and Prevention of Diabetic Foot Disease in Hospitals*. Diabetes UK, London.

Diabetes UK (2012) *Putting Feet First: Integrated Diabetes Foot Care Pathway*. Diabetes UK; London

Diabetes UK (2013) *Fast Track for a Foot Attack: Reducing Amputations (February 2013)*. Diabetes UK, London. Available at: <http://bit.ly/1Q8g7Xs> (accessed 14.01.2016)

Fard AS, Esmaelzadch M, Larijani B (2007) Assessment and treatment of a diabetic foot ulcer. *Int J Clin Pract* **61**: 1931–8

Glasper A (2010) Can the NICE quality standards help drive NHS improvements?, *Br J Nurs* **19**: 982–3

Guidroz AM, Luce KW, Denison DR (2010) Intergratedchange:

creating synergy between leader and organizational development. *Industrial and Commercial Training* **42**: 151–5

Gunther S, Taub N, Rogers S, Baker R (2013) What aspects of primary care predict emergency admission rates? A cross section study. *BioMed Central Health Services Research* **13**: 1–7

Harms PD, Crede M (2011) Emotional Intelligence and Transformational and Transactional Leadership: A Meta-Analysis. *Journal of Leadership and Organizational Studies* **17**: 5–17

Hayes C (2009) Interprofessional capacity building in diabetic foot management. *Br J Nurs* **18**: 804–10

Holman N, Young RJ, Jeffcoate WJ (2012) Variation in the recorded incidence of amputation of the lower limb in England. *Diabetologia* **55**: 1919–25

Jeffcoate W, Rayman G (2011) New guidelines for the diabetic foot: Let's make it a giant leap forward. *The Diabetic Foot Journal* **14**: 111–4

Kerr M (2012) *Foot Care for People with Diabetes: The Economic Case for Change*. NHS Diabetes and Kidney Care, London. Available at: <http://bit.ly/1ZYaFca> (accessed 12.01.2016)

Kristensen T, Laudicella M, Ejersted, Street A (2010) Cost variation in diabetes care delivered in English hospitals. *Diabetes Med* **27**: 949–57

Mahaffey K, Stanisstreet D, Ford M et al (2012) Role of the diabetes inpatient specialist nurse in preventing hospital admission from A&E. *Journal of Diabetes Nursing* **16**: 57–62

McInnes AD (2012) Diabetic foot disease in the United Kingdom: about time to put feet first. *J Foot Ankle Res* **5**: 1–7

Mooney J (2013) Foot infection and diabetes mellitus. *Podiatry Now* **16**: 16–21

Moran JW, Brightman BK (2000) Leading organizational change. *Journal of Workplace Learning* **12**: 66–74

Moulton C (2013) Diabetic footcare: part 1 avoiding amputations. *Journal of Diabetes* **43**: 8–16

Murdoch J, Barnes R, Pooler J et al (2014) Question design in nurse-led and GP-led telephone triage for same-day appointment requests: a comparative investigation. *BMJ Open* **4**: 1–9

Nason GJ, Strapp H, Kiernan C et al (2012) The cost utility of a multi-disciplinary foot protection clinic (MDFPC) in an Irish hospital setting. *Ir J Med Sci* **182**: 41–5

National Diabetes Inpatient Audit (2010) *National Diabetes Inpatient Audit 2010*. Available at: <http://bit.ly/1RzzQBh> (accessed 12.01.2016)

National Diabetes Inpatient Audit (2012) *National Diabetes Inpatient Audit 2011*. Available at: <http://bit.ly/1N4E0Zv> (accessed 12.01.2016)

National Diabetes Inpatient Audit (2013) *National Diabetes Inpatient Audit 2011–2012*. Available at: <http://bit.ly/1N4E0Zv> (accessed 12.01.2016)

NICE (2011) *Diabetic Foot Problems: Inpatient Management of Diabetic Foot Problems: NICE Clinical Guideline 119*. NICE, London. Available at: <http://bit.ly/1Ppe4tA> (accessed 12.01.2016)

NICE (2015) *Diabetic Foot Problems: Prevention and Management: NG19*. NICE, London. Available at: <http://bit.ly/1NdG8mM> (accessed 12.01.2016)

North Mersey Diabetes Network (2011) *Leading Change Event Diabetes Foot Care Pathway: Aintree Racecourse*. (1), pp1-12

Pope C, Chalder M, Moore L, Salisbury C (2005) What do other local providers think of NHS walk-in centres? Results of a postal survey. *Public Health* **119**: 39–44

Pope C, Halford S, Turnbull J et al (2013) Using computer support systems in NHS emergency and urgent care: ethnographic study using normalisation process theory. *BMC Health Serv Res* **13**: 1–13

Roberts P, Newton V (2011) Assessment and management of diabetic foot ulcers. *Br J Community Nursing* **16**: 485–90

Royal College of Nursing (2013) *Pressures on A&E are Putting Patients at Risk*. Available at: <http://bit.ly/1Skj5cN> (accessed 14.01.2016)

Stang D, Munro W (2015) Making Prevention pay. *The Diabetic Foot Journal* **18**: 1–4

TriePodD-UK (2013) *Podiatry Competency Framework For Integrated Diabetic Foot Care – A User's Guide*. SB Communications Group, London

Turner B (2013) Putting Feet First: Ensuring integrated foot care service for people living with diabetes and AQP going forward. *The Diabetic Foot Journal* **16**: 88–91

Yarwood-Ross L, Randall S (2013) Managing A Patients Diabetic Foot Ulcer. *Primary Health Care* **23**: 16–20