

Use of hormone replacement therapy in women with diabetes

Satinder Bal

Introduction

Coronary heart disease (CHD) is the major cause of mortality in women with type 2 (non-insulin-dependent) diabetes mellitus. Hormone replacement therapy (HRT) has generally favourable effects on cardiovascular risk factors. In this article, the known benefits of HRT in non-diabetic women and the potential benefits in women with diabetes mellitus are discussed. The results of a questionnaire study conducted among postmenopausal women attending a diabetes clinic to ascertain the prescription rates of HRT in diabetic women are presented.

The average age of the menopause in women is 50.8 years. Classically the diagnosis is made retrospectively following 12 months of amenorrhoea, although many women seek medical advice before this time because of the onset of menopausal symptoms. Eighty per cent of women experience vasomotor symptoms, which can be distressing and embarrassing.

Several studies have demonstrated an increased usage of hormone replacement therapy (HRT) in the past 10 years, although uptake of HRT remains less than 15% in women aged 40–65 years (Wilkes at al, 1991).

Ryan et al (1992) studied compliance with HRT in a group of women attending for bone mineral densitometry. They found a compliance rate of 74% following medical advice to continue therapy which fell to 61% at 12 months. Fear of cancer often deterred patients from starting treatment, and bleeding and weight gain were the main reasons for discontinuing therapy.

The immediate benefit gained by the patient is symptomatic relief. Hot flushing is the most commonly reported menopausal symptom and these vasomotor symptoms respond particularly well to HRT. Longer-term benefits are seen on the skeletal and cardiovascular systems.

Osteoporosis

Osteoporosis is defined by the World Health Organization (WHO) as 'a disease

characterised by low bone mass and microarchitectural deterioration of bone tissue, leading to enhanced bone fragility and a consequent increase in fracture risk' (Report of a WHO Study Group, 1994). It is therefore an important health hazard in menopausal women. Bone loss occurs at a rate of up to 4% per year for the first 8 years after the menopause, therefore fractures are an increasingly common source of morbidity in this group of patients.

The bone-sparing properties of oestrogens have been recognised for decades. Most studies indicate that fracture rates may be halved in current or recent past users of oestrogens.

Cardiovascular system

The incidence of cardiovascular disease in women rises steadily following the menopause, approaching that of men by age 70. Coronary heart disease (CHD) remains the leading cause of death in women over 65 years of age.

There is evidence that HRT can reduce death and disability from CHD; however, it is important to stress that CHD is multifactorial in causation, and overall risk factor modification remains the cornerstone of primary and secondary prevention. Epidemiological studies suggest that cardiovascular morbidity and mortality may be reduced by as much as 50% in postmenopausal women taking unopposed oestrogen therapy.

ARTICLE POINTS

1 Hormone replacement therapy (HRT) is underprescribed in the UK.

2 HRT can have major beneficial effects on the skeletal and cardiovascular systems.

3 Women with diabetes mellitus have an increased mortality from cardiovascular disease

4 HRT appears to have a favourable effect on cardiovascular risk factors.

5 More research on HRT and diabetic women is needed.

KEY WORDS

- Diabetes mellitus
- Menopausal women
- Hormone replacement therapy
- Cardiovascular disease

Satinder Bal is Specialist Registrar at Dumfries and Galloway Royal Infirmary, Dumfries

PAGE POINTS

1 Postmenopausal diabetic women have a much greater risk of developing cardiovascular disease than non-diabetic women.

2 Several cardiovascular risk factors may coexist in postmenopausal diabetic women.

3 HRT has been shown to improve glycaemic control as measured by HbA_{1c} levels.

4 HRT has had favourable effects on lipid profiles in both non-diabetic and diabetic women.

Although there are limited data on the use of oestrogen and progesterone combinations, the findings so far are encouraging.

Type 2 diabetes and risk of cardiovascular disease

Postmenopausal women with type 2 diabetes have a much greater risk of developing cardiovascular disease than non-diabetic women. The potential benefits of HRT in this group can only be extrapolated from results of studies in non-diabetic women as there is currently a paucity of research on HRT in women with diabetes mellitus. HRT has a favourable influence on several cardiovascular risk factors.

Insulin resistance

Reduced insulin sensitivity is a key factor in the clustering of cardiovascular risk factors observed in postmenopausal women with type 2 diabetes. Hyperlipidaemia, glucose intolerance, hypertension, impaired fibrinolysis and abdominal obesity often coexist in these individuals.

The PEPI trial was the first major, randomised, placebo-controlled trial examining the effects of HRT on heart disease risk factors in non-diabetic women (The Writing Group for the PEPI Trial, 1995). This trial demonstrated that patients receiving active treatment had slightly lower fasting insulin and glucose levels, although their 2-hour post-challenge glucose levels were slightly increased relative to the placebo group.

Brussard et al (1995) examined the effects of HRT in women with type 2 (non-insulin-dependent) diabetes mellitus. They employed the euglycaemic hyperinsulinaemic glucose clamp technique and demonstrated that HRT significantly reduced hepatic glucose production and improved glycaemic control as measured by the glycated haemoglobin (HbA_{1c}) level. It is possible that some of this beneficial effect may be offset by the addition of progestogens, although further studies are required to evaluate the effects of different HRT preparations upon insulin sensitivity in women with type 2 diabetes.

Dyslipidaemia

Patients with type 2 diabetes, particularly

females, have elevated plasma triglyceride levels and lower high-density lipoprotein (HDL) cholesterol concentrations, both of which are independent risk factors for CHD. Most studies examining the effects of HRT on lipid profiles have been undertaken in non-diabetic women, and have demonstrated favourable changes, i.e. a rise in HDL cholesterol concentration and a fall in low-density lipoprotein (LDL) cholesterol concentration.

Brussard et al (1995) demonstrated this rise in HDL cholesterol in a group of women with type 2 diabetes treated with oral oestradiol. The PEPI trial, however, found that unopposed oestrogen had a more favourable effect on HDL cholesterol than oestrogen given with continuous or cyclic synthetic progestogen. There were also differences depending on the progestogen: women treated with oestrogen and micronised progestogen had significantly higher cholesterol levels than women treated with oestrogen and medroxyprogesterone acetate.

Risks of treatment

Breast cancer

Fear of breast cancer is one of the main reasons women choose never to start HRT. The greatest risk factors for developing breast cancer are a strong family history of breast cancer and a biopsy indicating pre-malignant change. Breast cancer kills more young women than CHD, although women are five times more likely to die of an ischaemic cardiac event.

A large number of case-control and cohort studies have studied HRT use and breast cancer risk. These have produced conflicting results. There have been several meta-analyses of these studies which were largely biased towards oestrogen-only HRT, although more recent reviews have included combined preparations.

Hunt et al (1987) compared long-term oestrogen-only HRT and oestrogen-progestogen combinations and found a similar risk of breast cancer with the two preparations.

The meta-analysis by Colditz et al (1993) found that use of a combined preparation was associated with a relative risk of 1.2 at 5 years, which rose to 1.46 after 5–10

Table 1. Reasons why patients were not receiving HRT (n=65)

	Number	%
Never discussed treatment	50	77
Declined treatment	8	12
Valid contraindication	5	8
Diabetes deemed a contraindication	2	3

diabetes and 64 (72%) had type 2 diabetes. Overall, 24 (27%) were receiving HRT: 11 of the 25 patients (44%) with type 1 diabetes but only 13 of the 64 (20%) with type 2 diabetes. Sixty-five (73%) were not taking HRT, for a variety of reasons (Table 1). Significantly, 77% of patients had never discussed this form of treatment with a health-care professional.

The cardiovascular risk profiles of our cohort were examined. Fifty-one (57%) had documented coronary heart disease (CHD) or more than one risk factor for CHD in addition to diabetes, namely hyperlipidaemia (cholesterol >6.0mmol/litre, triglycerides >3.0mmol/litre, or on lipid-lowering therapy), hypertension, a history of smoking, or a family history of CHD. Only 25% of patients in this group were receiving treatment, although a small number (8%) had been offered treatment and declined.

Although this is a small survey, the results are interesting. The overall prescription rate of HRT was found to be almost double that in the general population. This difference may be explained partly by our very active local menopause clinic and heightened awareness of local GPs. We are currently planning to extend this survey to other diabetes clinics throughout the West of Scotland in order to increase the sample size and to assess geographical differences in prescription rates.

Conclusion

There is considerable evidence to suggest that the prescription of HRT can improve mortality in non-diabetic women. It should also reduce the burden of CHD in women with type 2 diabetes through a favourable effect on cardiovascular risk factors. Unless specific contraindications exist, these

women may achieve even greater benefits from treatment than their non-diabetic counterparts.

There is, however, a paucity of research on this topic. Further studies on the effects of the various forms of HRT in women with diabetes mellitus, who are a very high risk population with respect to vascular disease, are needed urgently. ■

Bal SK, McCrimmon RJ, Walker D, Frier BM (1997) The use of hormone replacement therapy in women with diabetes mellitus. *Diabetic Medicine* **14**: 55 A110 (Abstract)

Brussard HE, Gevers Leuven JA, Kluff C, Krass HMG (1995) Oestrogen replacement improves insulin response, lipoproteins and fibrinolysis in women with NIDDM. *Diabetologia* **38**: A25 (Abstract)

Colditz GA, Egan KM, Stampfer MJ (1993) Hormone replacement therapy and risk of breast cancer: results from epidemiological studies. *American Journal of Obstetrics and Gynecology* **168**(5): 1473–80

Fehér MD, Isaacs AJ (1996) Is hormone replacement therapy prescribed for postmenopausal diabetic women? *British Journal of Clinical Practice* **50**(8): 431–2

Hunt K, Vessey M, McPherson K et al (1987) Long term surveillance of mortality and cancer incidence in women receiving hormone replacement therapy. *British Journal of Obstetrics and Gynaecology* **94**: 620–35

Report of a WHO Study Group (1994) *Assessment of Fracture Risk and Its Application to Screening for Postmenopausal Osteoporosis*. WHO Technical Report Series 843. WHO, Geneva

Ryan PJ, Harrison R, Blake GH, Fogelman I (1992) Compliance with hormone replacement therapy (HRT) after screening for postmenopausal osteoporosis. *British Journal of Obstetrics and Gynaecology* **99**: 325–8

Sturdee DW, Barlow DH, Ulrich LG et al (1994) Is timing of withdrawal bleeding a guide to endometrial safety during sequential oestrogen-progestogen replacement therapy? *Lancet* **343**: 979–82

The Writing Group for the PEPI trial (1995) Effects of oestrogen or oestrogen/progestogen regimens on heart disease risk factors in postmenopausal women. *Journal of the American Medical Association* **273**: 199–208

Voight LF, Weiss NS, Chu J et al (1991) Progestogen supplementation of exogenous oestrogens and risk of endometrial cancer. *Lancet* **338**: 274–7

Wilkes HC, Meade TW (1991) Hormone replacement therapy in general practice: a survey of doctors in the MRC's General Practice Framework. *British Medical Journal* **302**: 1317–20

PAGE POINTS

1 HRT is generally considered to be grossly underprescribed in diabetic women.

2 However, the HRT prescription rate in this small study of diabetic women was double that in the general population.

3 Three-quarters of women in the study had never discussed HRT with a health professional.

4 Although more than half had CHD or more than one risk factor in addition to their diabetes, only a quarter were receiving treatment.

5 Diabetic women may achieve even greater benefits from HRT than non-diabetic women.