

Dietary carbohydrate restriction as a management strategy for adults with type 2 diabetes: Exploring the opinions of dietitians

Rosemary Huntriss, Ruth Boocock, Paul McArdle

Carbohydrate restriction is recognised as an appropriate dietary approach for people with type 2 diabetes. The aim of this study was to explore the opinions and practices of UK Registered Dietitians regarding this dietary approach. Data were collected by electronic questionnaires from 33 participants and underwent summative content analysis. Most respondents advise a moderate-carbohydrate diet to their patients, and the majority suggested that the high-carbohydrate intake promoted in public health guidance is inappropriate for adults with type 2 diabetes. A low-carbohydrate diet is offered by many dietitians and is considered achievable but not suitable for all. Importance was placed on individualisation of care.

There is a lack of evidence to recommend a single dietary approach for people with type 2 diabetes (Dyson et al, 2018); however, there remains significant interest in low-carbohydrate diets. There is some variation in practice amongst dietitians' dietary advice to adults with type 2 diabetes with respect to restricting carbohydrate (McArdle et al, 2017), despite nutrition guidelines including low-carbohydrate approaches and recent systematic reviews and meta-analyses supporting their use in the short term (Dyson et al, 2018; Huntriss et al, 2018; Sainsbury et al, 2018; McArdle et al, 2019). Evidence currently supports low-carbohydrate diets (<130 g per day) as an effective strategy for improving HbA_{1c} up to 12 months (Huntriss et al, 2018), but there is insufficient evidence to promote their effectiveness over other dietary approaches in the longer term (Sainsbury et al, 2018; McArdle et al, 2019).

UK nutrition guidelines for people with diabetes emphasise an individualised approach to dietary management, with low-carbohydrate diets being recommended as an option since 2011 (Dyson et al,

2011; 2018). A recent position statement from the British Dietetic Association (2018) draws the same conclusions.

The aim of this study was to explore the views and practices of UK Registered Dietitians (RDs) regarding dietary carbohydrate restriction as a management strategy for adults with type 2 diabetes. The results will further inform the debate within the dietetics profession and beyond concerning the appropriateness of this dietary intervention, and will provide insight as to how patients can be best supported to follow a low-carbohydrate diet.

Methods

This qualitative study used electronic questionnaires to collect data from RDs with clinical experience in type 2 diabetes management. Data were analysed using summative content analysis (Hsieh and Shannon, 2005). Bradford Teaching Hospitals NHS Foundation Trust acted as a sponsor for this study. Health Research Authority (HRA) approval was granted for this study (REC reference: 18/HRA/0014). Further ethical approval was not required for this study.

Citation: Huntriss R, Boocock R, McArdle P (2019) Dietary carbohydrate restriction as a management strategy for adults with type 2 diabetes: Exploring the opinions of dietitians. *Journal of Diabetes Nursing* 23: JDN104

Article points


1. Most of the dietitians surveyed in this study advise their patients with type 2 diabetes a moderate-carbohydrate approach, and the majority felt a low-carbohydrate diet was achievable for the right individuals with the right support.
2. Most also felt that the Public Health England guidance of a 50% energy intake from carbohydrates is inappropriate for this patient group.

Key words

- Diet and lifestyle
- Dietitian
- Low-carbohydrate diet
- Type 2 diabetes

Authors

Rosemary Huntriss is Specialist Diabetes Dietitian at Bradford Teaching Hospitals NHS Foundation Trust and Oviva UK Ltd; Ruth Boocock is Highly Specialist Diabetes Dietitian at Bradford Teaching Hospitals NHS Foundation Trust and Senior Lecturer in Dietetics at Teesside University; Paul McArdle is Lead Clinical Dietitian and Deputy Head of Nutrition at Birmingham Community Healthcare NHS Foundation Trust.



Diabetes UK nutrition guidelines for the prevention and management of diabetes. Available at: <https://bit.ly/2xR60mp>

Subjects and sampling

Convenience sampling was used to access participants for this study. Eligible participants were RDs with clinical experience in managing adults with type 2 diabetes. Those who were ineligible included dietitians not registered to practise in the UK and RDs who only offered general dietary advice to adults with type 2 diabetes in group settings. RDs who were employed by the study sponsor (Bradford Teaching Hospitals NHS Foundation Trust) were also excluded.

Study invitations were sent to 340 members of the British Dietetic Association's Specialist Diabetes Group via email. Participants confirmed that they met the eligibility criteria and gave their voluntary and informed consent to participate in the study prior to questionnaire completion. No incentives were offered to participants.

Survey design and administration

A pilot questionnaire was designed based on discussions in current literature (van Wyk et al, 2016; McArdle et al, 2017; Dyson et al, 2018; Huntriss et al, 2018), and sent to Specialist Diabetes Dietitians at Bradford Teaching Hospitals NHS Foundation Trust. Following a review of the process and questions, a final version of the questionnaire was agreed by the authors. The final version of the questionnaire was sent to participants via SurveyMonkey in May 2017.

The open-ended questions were developed with no pre-judgement and there were no leading questions. Participants could respond in free text to the questions. The questionnaire was available to complete for three weeks, as was communicated in the participant information sheet. No data were collected after this point.

Data analysis

Data were analysed using summative content analysis. This involves counting and comparing keywords or content (Hsieh et al, 2005). This act of assigning numerical values to a set of data may be used to quantify both manifest and latent content. Manifest content is evidence that is directly seen, such as the words written down in response to questions asked. Latent content refers to the underlying meaning of content, such as the interpretation of the answers to the questionnaire.

This methodology is commonly used to analyse open-ended survey questions (McKenna et al, 2017).

Each completed questionnaire was read through several times to allow the lead researcher (RH) to become familiar with the data. All questionnaires were printed out. Each question was analysed in turn. Codes from each question were identified and highlighted. Once codes from a question had been identified from all questionnaires, this process was repeated to ensure no codes were missed. Codes within each question were then counted and underlying context was interpreted through the generation of categories and themes. This process was done for each question individually. A second researcher (RB) independently analysed the codes for each question and then determined categories and themes. The codes, categories and themes of each researcher were compared, and a consensus was agreed. Relevant quotations were extracted to promote the authenticity and trustworthiness of the data (Elo and Kyngäs, 2008).

Results

Three hundred and forty dietitians were invited to take part in the study. Of these, 33 (9.7%) met the eligibility criteria, gave informed consent and participated. The answers to each question are described below.

Q1. What quantity of carbohydrate do you currently advise to adults with type 2 diabetes?

Thirty-three participants (100%) responded to this question. The frequency of responses can be found in *Table 1*.

Quantity

Nineteen participants (57.6%) responded to this question quoting grams per day. Five participants (15.2%) responded to this question by using percentage of total energy intake (% TEI).

Within these responses, 19 participants stated they might advise a moderate-carbohydrate diet (57.6%), ten a low-carbohydrate diet (30.3%) and five a high-carbohydrate diet (18.2%). Sixteen respondents presented a range, so not all respondents were exclusive to one category.

Educational messages

Common educational messages included advising a

reduction in carbohydrate intake ($n=5$) and filling one quarter of the plate with carbohydrate ($n=4$).

Individualisation

Several participants placed a strong emphasis on individualisation of care, suggesting their advice would depend on a number of factors, including the person, current intake, glycaemic control, weight, motivation, ability to make changes, medication, activity level and patient preference.

Q2. With regard to carbohydrate, how appropriate do you feel that the Eatwell Guide is in advising adults with type 2 diabetes (50% energy from carbohydrates)?

Thirty-three participants (100%) responded to this question.

Appropriateness of the Eatwell Guide for adults with type 2 diabetes

Sixteen participants responded that 50% TEI from carbohydrates, as promoted by the Eatwell Guide (Public Health England, 2016), is too high for adults with type 2 diabetes. An additional eight responded that the Eatwell Guide carbohydrate suggestion was inappropriate for adults with type 2 diabetes. Therefore, a total of 24 participants (72.7%) reported that the carbohydrate intake advised in the Eatwell Guide is either inappropriate or too high for adults with type 2 diabetes:

“[Type 2 diabetes] could be considered a condition of carbohydrate intolerance. We would not suggest to someone with coeliac disease to eat gluten, so why does the Eatwell Guide suggest to someone with [type 2 diabetes] to ensure their diet comprises 50% carbohydrate?”
(Participant 9)

One respondent commented that the Eatwell Guide does not echo the stance of Diabetes UK that there is little evidence for an ideal proportionate macronutrient intake for people with type 2 diabetes. Six respondents commented that the Eatwell Guide does not individualise advice, something considered important by this cohort.

Four respondents highlighted the difficulty that this nutritional model presents to overweight adults with type 2 diabetes or those who have insulin resistance.

Table 1. Frequency of responses to question 1.

Question: What quantity of carbohydrate do you currently advise to adults with type 2 diabetes?	
Response	Frequency count
Quantity advised	
Low-carbohydrate diet (<130 g/day or <26% TEI)	10
Moderate-carbohydrate diet (130–225 g/day or 26–45% TEI)	19
High-carbohydrate diet (>225 g/day or >45% TEI)	6
Educational messages	
Reduce carbohydrate intake	5
Allocate a quarter of the plate to carbohydrate	4
“Moderate” carbohydrate intake	2
Low-glycaemic-index foods	2
Eatwell Guide	1
Individualisation of care	
Dependent on factors (e.g. activity levels)	8
Carbohydrate intake should be individualised	7
Advice varies	4
No specific amount recommended	3

TEI=total energy intake.

The Eatwell Guide as a public health tool

Two participants recognised that the Eatwell Guide is a public health tool as opposed to being designed specifically for adults with type 2 diabetes. One participant suggested that it is suitable for healthy people and not those with a chronic condition such as type 2 diabetes. It was suggested that the Eatwell Guide is a useful tool to illustrate health messages such as the promotion of fruit and vegetable intake and reduction of high-sugar foods and drinks. However, one respondent suggested that 50% TEI from carbohydrates is likely to be too high even for the general population.

Trustworthiness of the Eatwell Guide

Two respondents discussed how, as it is a widely used tool produced by Public Health England, the Eatwell Guide is trusted by both patients and clinicians. One commented that the Guide can

Table 2. Frequency of responses to question 3.

Question: For adults with type 2 diabetes, what do you think the benefits of reducing dietary carbohydrate would be?

Response	Frequency count
Improved glycaemic control	29
Weight loss	24
Reduced requirement for/delayed progression onto medication	12
Reduced insulin resistance/circulating insulin	6
Reduced risk of diabetes-related complications	2
Improved cardiovascular outcomes	2
Decreased hunger	2
Reduced liver fat	1

be a starting point for healthcare professionals, but the other stated that patients might not want to decrease carbohydrate intake if they trust the recommendations of the Eatwell Guide.

What guidance would be appropriate for adults with type 2 diabetes?

Several respondents made suggestions as to how the Eatwell Guide could be adapted for adults with type 2 diabetes. Suggestions included reducing the TEI from carbohydrate to 25%, 40% or 25–50%, with one respondent suggesting the appropriateness of 25% TEI from carbohydrates as depicted on diet plates. Others suggested placing more of an emphasis on vegetables, salad and protein, and less on starchy carbohydrates.

Q3. For adults with type 2 diabetes, what do you think the benefits of reducing dietary carbohydrate would be?

Thirty-three participants (100%) responded to this question. The frequency of responses can be found in *Table 2*.

Glycaemic control

Twenty-nine participants (87.9%) reported that a reduction in dietary carbohydrate would improve glycaemic control in an adult with type 2 diabetes. Additionally or independently, several respondents identified individual elements of

glycaemic control that would be improved: blood glucose levels, HbA_{1c} and postprandial blood glucose levels.

Nine respondents reported that a reduction in medication would be seen and three reported that the progression onto diabetes medications would be delayed.

Weight loss

Twenty-four respondents (72.7%) attributed weight loss to a reduction in dietary carbohydrate. Two respondents acknowledged the greater satiating properties of the other macronutrients.

Other medical benefits

Other identified advantages of reducing dietary carbohydrate included decreased liver fat, delayed progression or reduced risk of diabetes-related complications, and improved cardiovascular outcomes such as lipid levels and blood pressure.

Q4. For adults with type 2 diabetes, what do you think would be the disadvantages of reducing dietary carbohydrate?

Thirty-two participants (97.0%) responded to this question.

Nutrition

Commonly reported concerns of reducing dietary carbohydrate were lack of fibre (*n*=9), increased fat intake (*n*=9), increased protein intake (*n*=6) and increased calorie intake (*n*=5). Nutritional deficiencies were stated as a concern (*n*=4), as was an insufficient intake of vitamins and minerals (*n*=4).

Physiology

The risk of hypoglycaemia was a common concern (*n*=7). Other mentioned disadvantages were lethargy (*n*=6), hunger (*n*=3), decreased concentration (*n*=2) and stress (*n*=1).

Practicality

Practical disadvantages, such as the implications of meal planning (*n*=4) and increased cost (*n*=2), were suggested. One participant responded that patients could find this difficult to follow, and another highlighted that patients may simply be unwilling to follow this diet. One respondent suggested that advising patients to reduce dietary carbohydrate

could cause confusion with regard to previous dietary advice (e.g. public health advice).

Three respondents raised questions about adherence to a reduced-carbohydrate diet.

No disadvantages

Three participants felt that, if managed appropriately, there would be no disadvantages to following a reduced-carbohydrate diet. One participant suggested setting a lower acceptable limit of carbohydrate intake. One participant did not respond to the question.

Q5. Do you think a low-carbohydrate diet (<130 g/day or <26% of total energy intake) is achievable for adults with type 2 diabetes? Please explain your answer

Thirty-three participants (100%) responded to this question. The frequency of responses can be found in *Table 3*.

Achievability

Twenty-eight participants (84.8%) felt that a low-carbohydrate diet is achievable for adults with type 2 diabetes, but there was a caveat attached to many responses, to include suitability, motivation and the need for support.

Barriers

Five participants (15.2%) felt that a low-carbohydrate diet would not be achievable for adults with type 2 diabetes, again offering some context.

Q6. How do you think patients can best be supported to follow a low-carbohydrate diet?

Thirty-two participants (97%) responded to this question. Support was categorised into three broad headings: personal support, group education and practical support, to include meal and snack ideas, recipes, books and apps. The frequency of the various responses can be found in *Table 4*.

Health messages

Three respondents highlighted the importance of the alignment of messages from healthcare professionals regarding this topic. The need for consistency in messages was highlighted in addition to the need for well-informed and robust guidelines to guide healthcare professionals.

Table 3. Frequency of responses to question 5.

Question: Do you think a low-carbohydrate diet (<130 g/day or <26% total energy intake) is achievable for adults with type 2 diabetes? Please explain your answer.

Response	Frequency count
Yes	28
Not for all with type 2 diabetes	7
Support or guidance needed	7
If motivated	7
Depends on the individual	4
Good understanding of nutrition required	3
If willing to try new foods	2
Cooking skills required	2
No	5
Cost is a barrier	4
Question of sustainability	4

“Twenty-eight participants (84.8%) felt that a low-carbohydrate diet is achievable for adults with type 2 diabetes, but there was a caveat attached to many responses, to include suitability, motivation and the need for support.”

Four respondents stated it is important to advise low-carbohydrate eating within the context of a healthy diet.

Q7. Please share any other comments or views you have regarding dietary carbohydrate restriction within this patient group

Twenty participants (60.6%) responded to this question. The quotes below summarise the points made:

“The low-fat healthy eating diet has led to an increase in portion sizes of rice/pasta/potato and an attitude that these are healthy, so the more the better – this needs to be changed, and talking about reducing carbohydrate and explaining why really helps.” (Participant 30)

“Evidence suggests weight loss is key to management of [type 2 diabetes] but no evidence as to the best approach. Carb reduction not only offers



Read more online

How to improve carbohydrate awareness

In the ongoing How to... series of quick reference guides for primary care professionals, Jane Diggle provides all the background information and practical tips needed to discuss carbohydrates with patients.

Diabetes & Primary Care **21**: 85–6

[Click here to access](#)

Table 4. Frequency of responses to question 6.

Question: How do you think patients can best be supported to follow a low-carbohydrate diet?

Response	Frequency count
Personal support	
Regular support	7
Dietetic support	6
One-to-one	4
Group	4
Peer support	3
Diabetes professional	2
Social media	2
Eatwell Guide	1
Education	
Carbohydrate awareness	10
Structured education	2
Practical support	
Ideas for meals and snacks	16
Recipes	7
Apps	5
Written literature	5
Books	4
Help with portion control	4
Diabetes medication management	3
Blood glucose monitoring	1

opportunities for weight loss but reduces blood glucose levels as an added bonus.” (Participant 27)

“I think it would definitely be beneficial for diabetes management however the majority would struggle to achieve a low CHO diet. General CHO awareness is of importance.” (Participant 32)

“People with non-diabetic hyperglycaemia/ pre-diabetes/impaired fasting glucose, however it is labelled, should be strongly advised to reduce their carbohydrate intake.” (Participant 32)

“Due to lack of long-term evidence regarding safety of low-carbohydrate diets (especially those that recommend 50–90 g per day) I am not comfortable recommending them for long term.” (Participant 7)

“Advice HAS to be individualised.” (Participant 13)

“Would like more research and evidence to be available as to benefits of this type of diet and sustainability long-term.” (Participant 25)

“Would value more support and resources from the BDA/Diabetes UK, etc., to support patients who wish to follow a low carbohydrate diet.” (Participant 26)

Discussion

This study used summative content analysis to explore the opinions of UK RDs regarding carbohydrate restriction as a management strategy in adults with type 2 diabetes. Thirty-three UK RDs from a diabetes specialist interest group took part. Participants were asked six focused questions and a final broad question inviting other comments on the topic.

The participants were overwhelmingly in favour of advising a restricted-carbohydrate diet to adults with type 2 diabetes, with 29 reporting they may advise low- or moderate-carbohydrate diets to adults with type 2 diabetes. Importantly, there was a strong emphasis around individualisation and there was acknowledgement that there is no one-size-fits-all approach.

A significant number of participants (n=24) also believe the current Public Health England dietary guidance, the Eatwell Guide, is either too high in carbohydrate or inappropriate for adults with type 2 diabetes. Some participants highlighted its purpose as a public health tool as opposed to one designed to support those with specific health conditions. Consequently, these data suggest that the RDs sampled believe the Eatwell Guide, and the message that 50% TEI should come from carbohydrates, should be interpreted with caution when advising adults with type 2 diabetes.

Most participants felt that a low-carbohydrate diet is achievable for adults with type 2 diabetes, but this

conclusion requires further interpretation. Although participants felt it could be achieved, several raised important points and concluded that this diet would not be suitable for all adults with type 2 diabetes. Namely, an individual would likely need to be motivated and would likely need support. For RDs who felt that this diet is not achievable, cost and sustainability were the two key themes of concern, the latter of which is acknowledged in the published literature (van Wyk et al, 2016).

Participants were largely in agreement with respect to the perceived benefits of carbohydrate restriction, as almost all expected an improvement in glycaemic control with intentional weight loss where appropriate. A greater divergence of views was observed with regard to the possible disadvantages of reducing dietary carbohydrate. The overarching themes were potential nutritional concerns, health-related implications and practicality, with significant variation within the themes; the most commonly suggested disadvantages were lack of fibre, increased fat intake and increased risk of hypoglycaemia. Regular support, dietetic support and practical ideas for meals, snacks and recipes were cited by respondents as the key support needed by patients to follow a low-carbohydrate diet. Carbohydrate awareness was mentioned; however, despite being popular, this remains a poorly defined term among dietitians (McArdle et al, 2017).

Study strengths and limitations

Strengths of this research include the sampling from a population of RDs interested in diabetes through existing networks, and the rigorous analysis of the data using triangulation by a second investigator.

An obvious limitation to this study was the low response rate. Internet-based surveys are known to be time- and cost-efficient but can also have drawbacks such as selection bias and poor response rates (Fan and Yan, 2010). Sampling and sample size are important considerations for all research, and in a qualitative study such as this, where the response rate cannot be predicted, the validity and reliability of the data must be judged on factors such as saturation and replication of responses across themes (Elo et al, 2014). There was frequent replication and consistency of responses within themes in many questions, and these were identified by both authors involved in the data analysis, indicating

the data were sufficiently saturated for these results to be reliable and trustworthy. The frequencies of all responses were reported in tables to promote transparency of the results.

Qualitative research does not claim generalisability, and these results are an exploration of the views of a UK-wide group of RDs with a special interest in diabetes. As previously referenced, online surveys can be subject to selection bias, and the relevance of the topic to the responders is also a major known factor in predicting the response rate for any survey (Dillman and Smyth, 2007). As a result, we are not claiming generalisability of these results but sharing the opinions of those clinicians who took part in the study. Equally, the study gave the opportunity for those who may have been against carbohydrate restriction to share their opinions, so we feel confident that the findings are valid. Furthermore, there were no incentives used in this study, a technique known to increase response rate (Fan and Yan, 2010).

Conclusions

In conclusion, UK RDs hold generally positive views of carbohydrate restriction as a management strategy for adults with type 2 diabetes, and their perceived benefits match the evidence base. There remains variation in clinical practice, but many RDs feel that UK public health guidance recommending a high-carbohydrate diet may not be suitable for adults with type 2 diabetes. Furthermore, consideration may need to be given as to how RDs may be supported in the provision of evidence-based low-carbohydrate dietary advice to adults with type 2 diabetes who choose to follow this approach over other dietary strategies, in a nutritionally balanced, acceptable and sustainable manner. The results of this study may not be generalisable but show important data regarding the opinions and practices of RDs supporting patients with type 2 diabetes in the dietary management of carbohydrate. ■

Acknowledgements

The authors would like to thank the sponsor of the study, Bradford Teaching Hospitals NHS Foundation Trust. In particular, we would like to thank the Head of Nutrition and Dietetics and Specialist Diabetes Dietitians. We would also like to extend our thanks to Julie Taplin and the BDA Diabetes Specialist Group.



Read more
online

Low-carbohydrate diets for children and young people with type 1 diabetes: Unpicking the evidence

A detailed review of the current evidence regarding low-carbohydrate diets and the potential risks associated with their use in young people, as well as advice on supporting and monitoring those patients who choose to follow this dietary approach.

Journal of Diabetes Nursing
23: JDN095

[Click here to access](#)

“UK Registered Dietitians hold generally positive views of carbohydrate restriction as a management strategy for adults with type 2 diabetes, and their perceived benefits match the evidence base.”

Author contributions

RH was responsible for protocol development, study design and conduct, data collection, analysis and interpretation, and manuscript writing. RB was responsible for review of protocol and study design, validity checking of data analysis and interpretation, and manuscript writing. PM was responsible for review of protocol and study design, and manuscript writing.

Conflict of interest

The authors declare no conflict of interest.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

British Dietetic Association (2018) *Policy statement: Low carbohydrate diets for the management of type 2 diabetes in adults*. BDA, London. Available at: <https://bit.ly/35CkdD6> (accessed 20.10.19)

Dillman DA, Smyth JD (2007) Design effects in the transition to web-based surveys. *Am J Prev Med* **32**(5 Suppl): S90–6

Dyson PA, Kelly T, Deakin T et al; Diabetes UK Nutrition Working Group (2011) Diabetes UK evidence-based nutrition guidelines for the prevention and management of diabetes. *Diabet Med* **28**: 1282–8

Dyson PA, Twenefour D, Breen C et al (2018) Diabetes UK evidence-based nutrition guidelines for the prevention and management of diabetes. *Diabet Med* **35**: 541–7

Elo S, Kyngäs H (2008) The qualitative content analysis process. *J Adv Nurs* **62**: 107–15

Elo S, Kääriäinen M, Kanste O et al (2014) Qualitative content analysis: a focus on trustworthiness. *SAGE Open*. DOI: 10.1177/2158244014522633

Fan W, Yan Z (2010) Factors affecting response rates of the web survey: a systematic review. *Comput Hum Behav* **26**: 132–9

Hsieh HF, Shannon SE (2005) Three approaches to qualitative content analysis. *Qual Health Res* **15**: 1277–88

Huntriss R, Campbell M, Bedwell C (2018) The interpretation and effect of a low-carbohydrate diet in the management of type 2 diabetes: a systematic review and meta-analysis of randomised controlled trials. *Eur J Clin Nutr* **72**: 311–25

McArdle PD, Greenfield SM, Avery A et al (2017) Dietitians' practice in giving carbohydrate advice in the management of type 2 diabetes: a mixed methods study. *J Hum Nutr Diet* **30**: 385–93

McArdle PD, Greenfield SM, Rilstone SK et al (2019) Carbohydrate restriction for glycaemic control in type 2 diabetes: a systematic review and meta-analysis. *Diabet Med* **36**: 335–48

McKenna L, Brooks I, Vanderheide R (2017) Graduate entry nurses' initial perspectives on nursing: content analysis of open-ended survey questions. *Nurse Educ Today* **49**: 22–6

Public Health England (2016) *The Eatwell Guide*. PHE, London. Available at: <https://bit.ly/2Jo0YBB> (accessed 20.10.19)

Sainsbury E, Kizirian NV, Partridge SR et al (2018) Effect of dietary carbohydrate restriction on glycaemic control in adults with diabetes: a systematic review and meta-analysis. *Diabetes Res Clin Pract* **139**: 239–52

van Wyk HJ, Davis RE, Davies JS (2016) A critical review of low-carbohydrate diets in people with type 2 diabetes. *Diabet Med* **33**: 148–57