

The 'ABC' approach to preventing diabetic foot problems

Ranjna Garg, Anil Kumar Agarwal, Anita Barker, Vinod Patel

ARTICLE POINTS

1 Diabetes-related foot ulcers are common and have high associated mortality.

2 Peripheral neuropathy and peripheral vascular disease contributes to diabetic foot complications.

3 A multidisciplinary approach is the key issue for quality care.

4 Regular screening, glycaemic control and blood pressure control should be optimal.

5 Appropriate footwear and risk factor control can reduce foot complications in diabetes.

KEY WORDS

- Foot ulcer
- Lower extremity amputation
- Mnemonic tool
- Foot complications

Ranjna Garg is Specialist Registrar at Manor Hospital, Walsall; Anil Kumar Agarwal is Specialist Registrar at Good Hope Hospital, Sutton Coldfield; Anita Barker is Chief Specialist Podiatrist and Vinod Patel is Consultant Physician and Diabetologist at George Eliot Hospital, Nuneaton.

Introduction

Diabetes-related foot complications are the main cause of non-traumatic lower limb amputation (Dang and Boulton, 2003). A simple, logical approach can reduce the incidence of diabetes-related foot ulcers and amputations (Malone et al, 1989). Careful assessment, monitoring, control of risk factors and identification of the at-risk foot are important. In this article, the authors describe a simple strategy aimed at reducing the incidence of diabetes-related foot ulcers and amputations. This mnemonic tool is intended as a simple set of guidelines for professionals working in the assessment and management of the diabetic foot.

The provision of optimal foot care is of paramount importance in avoiding catastrophic foot complications from diabetes. According to Calle-Pascual et al (2001), a thirteen-fold reduction in the incidence of foot-related complications of diabetes could be achieved through the provision of optimum foot care.

The diabetes team at the George Eliot Hospital in Nuneaton has previously published a mnemonic template (the 'Alphabet Strategy') for providing comprehensive diabetes care (Patel and Morrissey, 2002). An audit (Lee et al, 2003) showed that use of the Alphabet Strategy resulted in better outcomes in terms of optimal systolic and diastolic blood pressures, eye and foot examination frequency, appropriate use of 'guardian' drugs (e.g. aspirin, angiotensin-converting enzyme [ACE] inhibitors, statins) and coronary heart disease risk scores. The strategy is easily adapted and applicable in primary care and secondary care as well as for healthcare providers in the community. It can also be used as an educational tool.

As a subset of the recommendations set out in the template, the authors present a simple yet effective 'ABC' approach to identify, prevent and minimise the risk of diabetic foot-related complications – mainly amputations.

The ABC approach was designed to provide a simple set of guidelines for professionals working in assessment, treatment and management of the diabetic foot.

The 'ABC' approach

'A' is for assessment, advice and access

Assessment

In people with diabetes, the lifetime risk of developing a foot ulcer is as high as 15% (Boulton, 2004). Lavery et al (1998) demonstrated that the risk of foot ulceration is higher in males and in patients with poor glycaemic control, with diabetes of 10 years or more, with the presence of deformity or prior amputation or with subjective evidence of peripheral neuropathy. Assessment is important in the detection of early neuropathy and ischaemia. Moulik et al (2003) reported the presence of peripheral neuropathy in 61% of patients presenting to the foot clinic for the first time with a foot ulcer.

Identifying the at-risk foot early is important. The presence of sensory loss (e.g. as assessed with a 10g monofilament), ischaemia (as identified by absent pulses and reduced ankle brachial pressure index), deformity, callus, joint instability or oedema should all be assessed as they help in identifying the at-risk foot (Reiber et al, 1999).

The risk of ulcer formation is high in those patients with a history of foot ulceration (Lavery et al, 1998). In addition, Pecoraro et al (1990) demonstrated that in more than 80% of cases, amputation was preceded by foot ulceration. Therefore, interventions to

reduce the risk of ulceration are important. Both the use of orthotic devices (Colagiuri et al, 1995) and pressure relief (Lavery et al, 1996) have been shown to reduce the risk of foot ulceration.

All diabetes patients should have regular foot assessment as part of their diabetic foot programme (Peters et al, 2001), which should include complete neurovascular assessment.

Advice

Advice is important to the prevention of diabetic foot complications. For example, advice on re-vascularisation should be sought from vascular surgeons by podiatrists/chiropractors when dealing with ischaemic limbs and/or ulcers. Furthermore, all people with diabetes should be given advice on appropriate foot care, daily foot inspection and avoidance of trauma. Advice on smoking cessation also should be given and reinforced at every opportunity.

Access

All diabetes patients should have access to a

trained podiatrist/chiropractor.

'B' is for blood pressure

The United Kingdom Prospective Diabetes Study (UKPDS; UKPDS Group, 1998) showed that diabetes-related microvascular and macrovascular complications are reduced when tight blood pressure control is achieved. Blood pressure targets should therefore be strictly followed in people with diabetes.

'C' is for control of infection

The presence of infection delays ulcer healing (Pecoraro et al, 1990). In a study by Armstrong et al (1998), patients with wound infection were at higher risk of amputation than people with non-infected foot ulcers. In the authors' opinion, the presence of infection should be diligently searched for in non-healing or recurrent foot ulcers. Aggressive and appropriate antibiotic therapy should therefore be instituted in those people with infected ulcers (Lipsky, 2004).

PAGE POINTS

1 All diabetes patients should have regular foot assessment as part of their diabetic foot programme, which should include neurovascular assessment.

2 Advice is important to the prevention of diabetic foot complications.

3 In the authors' opinion, the presence of infection should be diligently searched for in non-healing or recurrent foot ulcers.

Table 1. ABC approach tick-box proforma used in Nuneaton

A: Advice, assessment, access	
Foot inspection	<input type="checkbox"/>
Smoking cessation	<input type="checkbox"/>
Assessment	<input type="checkbox"/>
Need for re-vascularisation	<input type="checkbox"/>
Access to podiatrist/chiroprapist	<input type="checkbox"/>
B: Blood pressure optimal	
<input type="checkbox"/>	
C: Control of infection	
<input type="checkbox"/>	
D: Debridement, dressings	
Debridement	<input type="checkbox"/>
Dressings	<input type="checkbox"/>
E: Education	
<input type="checkbox"/>	
F: Footwear, Fluid retention	
Footwear needed (specify type)	<input type="checkbox"/>
Fluid retention	<input type="checkbox"/>
G: Guardian drugs	
Asprin	<input type="checkbox"/>
ACE inhibitor	<input type="checkbox"/>
Blood pressure lowering	<input type="checkbox"/>
Lipid lowering	<input type="checkbox"/>
H: Complete	
<input type="checkbox"/>	

'D' is for debridement and dressings

Debridement

Adequate debridement has shown improved healing in diabetic foot ulcers (Steed et al, 1996; Jeffcoate et al, 2004). Regular wound inspection and debridement cleanses the wound surface; removes dead tissue, callus, bone and pathogenic bacteria; drains pus; and promotes healing.

Callus formation is a consequence of chronic continued foot pressure and leads to ulcer formation (Boulton, 2004). Debridement in ischaemic or neuroischaemic ulcers should be limited to removal of slough/pus.

Dressings

Dressings maintain the moist wound environment, help avoid trauma and minimise the risk of infection. Specially formulated dressings may absorb exudates and maintain hydration status in the wound. Many types of dressings are available, such as non-adhesive paraffin-impregnated and allogenic skin substitutes (e.g. Dermagraft [Smith and Nephew, Hull], Graftskin [Novartis, Camberley], Regranex [Janssen-Cilag, High Wycombe]). Specialist advice should be sought from a podiatrist/chiroprapist in choosing the most suitable dressing.

'E' is for education

People with diabetes need education, re-education and reinforced education. Faglia et al (2001) demonstrated a reduction in new ulceration/amputation in people with diabetes that were given intensive education. In the authors' opinion, education should be an ongoing process and addressed at each clinic visit. Every opportunity should be taken to remember this aspect of care. Stress should be placed on daily inspection of the feet. Patients should be educated to contact the foot care team early.

Similarly, health professionals should be educated to recognise early warning signs and urgent action should be taken when problems are detected.

'F' is for footwear and fluid retention

Footwear

Skin breaks when circulation is compromised by pressure. In the presence of neuropathy, this process can remain unnoticed until an ulcer develops. Pressure relief, therefore, forms an integral part of optimal foot care. Pressure reduction and/or off-loading is central to foot ulcer healing (Armstrong and Lavery, 1998; Boulton, 2004).

When deformity is present (e.g. hammer toes, prominent metatarsal heads, bunions), correct footwear (e.g. wide-fitting shoes) can prevent an ulcer from appearing. People with Charcot foot need specially moulded shoes to avoid repetitive trauma. Methods of off-loading include bed rest, total contact casting, removable walking casts, surgical shoes, felted foam dressings and half-shoes. Total contact casting is considered the 'gold standard' modality for off-loading (Lavery et al, 1996).




Figure 1. Use of the ABC approach in Nuneaton. The authors suggest this protocol could also be useful in other areas.

Fluid retention

Fluid retention in lower extremities can cause fluid-filled blisters that rupture to leave a raw surface. Unrecognised raw surfaces could emerge as a nasty ulcer. Fluid retention in the lower limb should therefore be minimised by, for example, simple elevation of foot, judicious use of diuretics, treatment of heart failure, and TED (thromboembolic deterrent) stockings (American Diabetes Association [ADA], 1999).

'G' is for 'guardian' drugs

'Guardian' drugs is the authors' term for the specific drugs that are used in treatment of the metabolic syndrome. Examples are lipid-lowering therapies, anti-platelet agents and ACE inhibitors. These therapeutic agents should be properly initiated and doses titrated to achieve specific targets (e.g. blood pressure and cholesterol targets).

The quest for ideal glycaemic control should not be forgotten and every effort should be made to obtain an HbA_{1c} level of 7% or less in people with diabetes (ADA, 2004). This should be a continued process by use of drugs, diet and exercise.

Diabetes UK currently recommends aspirin use in people with diabetes who are

over 30 years of age and have additional risk factors (e.g. smoking, overweight, dyslipidaemia; Diabetes UK, 2001). In addition, current recommendations from the ADA are to use low-dose aspirin as a primary prevention strategy in people with type 2 diabetes who are above 40 years of age or who are at high risk of cardiovascular disease. Dosages as low as 75 mg/day are effective (Colwell and ADA, 2004).

Use of the strategy in practice

The ABC approach has been incorporated into the diabetes care pathways in the diabetes centre at Nuneaton. Each patient's clinic letters include the tool mnemonics and, during the patient's visit, the outpatient nurse and consultant physician ensure these are ticked. In this way, people who fail a parameter are easily identified and corrected – e.g. patients who are not taking aspirin when they should be or who are not meeting blood pressure targets.

The ABC strategy forms an initial assessment tool at the time of a patient's first clinic visit and is then used at annual review visits. It is for use by all healthcare professionals who provide ongoing care to people with

PAGE POINTS

1 The ABC strategy forms an initial assessment tool at the time of a patient's first clinic visit and is then used at annual review visits.

2 The ABC strategy proforma is followed and ticked by the outpatient or diabetes specialist nurse in the diabetes clinic and by the chiropodist or podiatrist in the foot clinic.

3 Any unticked boxes are addressed in the clinic by the consultant physician.

4 In the authors' experience, use of the ABC strategy in this way has improved outcomes.

PAGE POINTS

1 The ABC strategy is a mnemonic approach that highlights the various aspects diabetic foot care.

2 Use of the mnemonic has made it simple to use effectively by all care providers.

3 The results of audits to examine the effects of the Alphabet Strategy for diabetes care in general (of which this ABC approach to diabetic foot care is one component) suggested improved outcomes in patient care.

diabetic foot problems. In Nuneaton, the podiatrist and chiropodist use the ABC protocol regularly. See *Table 1* for a proforma template used in the Nuneaton clinical setting.

The proforma is followed and ticked by the outpatient or diabetes specialist nurse in the diabetes clinic and by the chiropodist or podiatrist in the foot clinic. Any unticked boxes are addressed in the clinic by the consultant physician. In this way, for example, anti-hypertensive therapy may be intensified, aspirin treatment initiated or the patient initiated on to insulin therapy if glycaemic control is sub-optimal.

In the authors' experience, use of the ABC strategy in this way (*Figure 1*) has improved outcomes in terms of:

- people with diabetes meeting blood pressure, lipid and glycaemic targets
- the early identification of people at high risk of foot ulceration or ulcer recurrence.

Conclusion: 'H' is for healing

The authors believe that if stress is placed on following points 'A' to 'G' in the care of diabetes patients then diabetic foot ulcers can be prevented, identified early when present and treated appropriately to be more effective for complete healing.

The ABC strategy is a mnemonic approach that highlights the various aspects of diabetic foot care. Use of the mnemonic has made it simple to use effectively by all care providers. The results of audits carried out in Nuneaton (e.g. Lee et al, 2003) to examine the effects of the Alphabet Strategy for diabetes care in general (of which this ABC approach to diabetic foot care is one component) suggested improved outcomes in patient care, including in relation to the diabetic foot. ■

American Diabetes Association (1999) Consensus Development Conference on Diabetic Foot Wound Care: 7-8 April 1999, Boston, Massachusetts. American Diabetes Association. *Diabetes Care* **22**(8): 1354-60

American Diabetes Association (2004) Standards of medical care in diabetes. *Diabetes Care* **27**(Suppl 1): S15-35

Armstrong DG, Lavery LA (1998) Evidence-based options for off-loading diabetic wounds. *Clinics in Podiatric Medicine and Surgery* **15**(1): 95-104

Armstrong DG, Lavery LA, Harkless LB (1998) Validation of a diabetic wound classification system. The contribution of depth, infection, and ischemia to risk of amputation. *Diabetes Care* **21**(5): 855-9

Boulton AJ (2004) The diabetic foot: from art to science. The 18th Camillo Golgi lecture. *Diabetologia* **47**(8): 1343-53

Calle-Pascual AL, Duran A, Benedi A, Calvo MI, Charro A, Diaz JA, Calle JR et al (2001) Reduction in foot ulcer incidence: relation to compliance with a prophylactic foot care program. *Diabetes Care* **24**(2): 405-7

Colagiuri S, Marsden LL, Naidu V, Taylor L (1995) The use of orthotic devices to correct plantar callus in people with diabetes. *Diabetes Research & Clinical Practice* **28**(1): 29-34

Colwell JA; American Diabetes Association (2004) Aspirin therapy in diabetes. *Diabetes Care* **27**(Suppl 1): S72-3

Dang CN, Boulton AJ (2003) Changing perspectives in diabetic foot ulcer management. *International Journal of Lower Extremity Wounds* **2**(1): 4-12

Diabetes UK (2001) *Care recommendation: Aspirin treatment in diabetes*. Diabetes UK, London

Faglia E, Favales F, Morabito A (2001) New ulceration, new major amputation, and survival rates in diabetic subjects hospitalized for foot ulceration from 1990 to 1993: a 6.5-year follow-up. *Diabetes Care* **24**(1): 78-83

Jeffcoate WJ, Price P, Harding KG; International Working Group on Wound Healing and Treatments for People with Diabetic Foot Ulcers (2004) Wound healing and treatments for people with diabetic foot ulcers. *Diabetes/Metabolism Research and Reviews* **20** (Suppl 1): S78-89

Lavery LA, Armstrong DG, Vela SA, Quebedeaux TL, Fleischli JG (1998) Practical criteria for screening patients at high risk for diabetic foot ulceration. *Archives of Internal Medicine* **158**(2): 157-62

Lavery LA, Vela SA, Lavery DC, Quebedeaux TL (1996) Reducing dynamic foot pressures in high-risk diabetic subjects with foot ulcerations. A comparison of treatments. *Diabetes Care* **19**(8): 818-21

Lee J, Sankaranarayan S, Shaikh S, Chellamuthu C, Morrissey J, Patel V; The Alphabet POEM Project (Practice of Evidence-Based Medicine) (2003) Systemic application of the Alphabet strategy improves diabetes care parameters and coronary artery disease risk in patients with type 2 diabetes mellitus. *Diabetes* **52**(S1): A522

Lipsky BA (2004) Medical treatment of diabetic foot infections. *Clinical Infectious Diseases* **39**(Suppl 2): S104-14

Malone JM, Snyder M, Anderson G, Bernhard VM, Holloway GA Jr, Bunt TJ (1989) Prevention of amputation by diabetic education. *American Journal of Surgery* **158**(6): 520-3

Moulik PK, Mtonga R, Gill GV (2003) Amputation and mortality in new-onset diabetic foot ulcers stratified by etiology. *Diabetes Care* **26**(2): 491-4

Patel V, Morrissey J (2002) The Alphabet Strategy: The ABC of reducing diabetes complications. *British Journal of Diabetes & Vascular Disease* **2**: 58-9

Pecoraro RE, Reiber GE, Burgess EM (1990) Pathways to diabetic limb amputation. Basis for prevention. *Diabetes Care* **13**(5): 513-21

Peters EJ, Lavery LA; International Working Group on the Diabetic Foot (2001) Effectiveness of the diabetic foot risk classification system of the International Working Group on the Diabetic Foot. *Diabetes Care* **24**(8): 1442-7

Reiber GE, Vileikyte L, Boyko EJ, del Aguila M, Smith DG, Lavery LA, Boulton AJ (1999) Causal pathways for incident lower-extremity ulcers in patients with diabetes from two settings. *Diabetes Care* **22**(1): 157-62

Steed DL, Donohoe D, Webster MW, Lindsley L (1996) Effect of extensive debridement and treatment on the healing of diabetic foot ulcers. Diabetic Ulcer Study Group. *Journal of the American College of Surgeons* **183**(1): 61-4

UK Prospective Diabetes Study Group (1998) Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. *British Medical Journal* **317**(7160): 703-13