

The “processed truth”



Kate Walker

Owner of Diabetes and You
and of Be Nice To You

In this short viewpoint piece, I provide an alternative look at the basis underlying dietary advice for people with type 2 diabetes and obesity.

Information in this area is seemingly conflicting, but it is common for obese people with type 2 diabetes to be advised to eat a low-fat diet. In my view, there is insufficient clinical evidence to support low-fat diets as the optimal approach (e.g. Alhazmi et al, 2012) and, in fact, there is more evidence to suggest eating a high-protein or Mediterranean diet is beneficial (Brown et al, 2010; Swedish Council on Health Technology Assessment, 2013). Are we, however, asking ourselves the wrong questions about diet and therefore getting the wrong answers? What answers would we get to the question “What are we ‘designed’ to eat to be healthy?”

The earliest fossil remains of *Homo sapiens* date back to approximately 200 000 years ago, and genetically our species is approximately 99.9% similar to *Homo erectus*, a close relative in the hominin family tree that has been dated back as far as 1.9 million years ago (Hines, 2009). For millennia our way of life and our food revolved around what we could grow, hunt and gather from nature (Leach et al, 2006).

Manufactured fats, chemical fertilisers, fortification, processed foods and genetic modification are all essentially modern phenomena, generally being from the past 100 years (Schmidt, 2005; Samaniego-Vaesken et al, 2012). Genetically, we have not changed over this short period, but our food has changed dramatically (Schmidt, 2005; Hines, 2009). There has been insufficient time for natural selection to drive the evolution of metabolic processes to which a typical westernised diet and lifestyle will not be detrimental (Coffey, 2001).

Each human is different and will have grown up, or be growing up, in a different

environment; as such, our macronutrient ratios requirements will vary (Price, 2003). Simple broad-brush approaches to adjusting macronutrient ratios with a goal of obtaining weight loss may, therefore, not be appropriate.

A very different way of thinking about nutrition is that we may lose, and therefore need to replace, up to 3 million cells a second (Mack, 2001). For this replacement of cells, the body requires adequate intake of nutrients, amino acids, trace elements and minerals. This brings me to my next point, which concerns processed foods.

Processed foods

Processed food products contain minimal “natural nourishment” on account of pasteurisation, homogenisation, fortification, the addition of numerous chemicals (to enhance taste and improve shelf-life) and re-heating (Webster et al, 2010; Van Gysel et al, 2012; NHS Choices, 2013; NutritionData.com, 2013). These processes can make the food more difficult to digest in the stomach (McMichael, 2005) and may have made the remaining nutrients more difficult to access (Fallon and Enig, 2001). Simply put, highly processed foods do not provide nourishment and calories in the optimal ratios for our body’s needs (Hockett and Haws, 2003).

Little research has been published on the overall impact of processed foods on obesity and type 2 diabetes. There is, however, evidence to show the impact on our bodies of common constituents of processed foods, such as high-fructose corn syrup (HFCS) and trans fats.

Effect of high-fructose corn syrup on the body

HFCS is often used as an alternative to sucrose and is found in most processed meals

(Howard and Wylie-Rosett, 2001), but it does not currently need to be labelled explicitly on packaging in the UK. Fructose is broken down differently from glucose and sucrose, and has to be processed and stored in the liver (Basaranoglu, 2013). Unlike glucose, fructose does not stimulate insulin secretion or enhance leptin production and, because insulin and leptin act as key afferent signals in the regulation of food intake and body weight, dietary fructose may contribute to increased energy intake and weight gain (Bray et al, 2004). Consumption of fructose is also associated with adverse alterations of plasma lipid profiles (Goran et al, 2013), metabolic changes (Basaranoglu, 2013), type 2 diabetes (Goran et al, 2013) and impaired cognitive function (Lakhan and Kirchgessner, 2013). These observations indicate a role of HFCS in the obesity epidemic, but more research is needed in this area before definitive conclusions can be drawn.

Effect of processed fats on the body

Compared with natural, unprocessed cis fatty acids, trans fats have a different effect on lipid profiles (Mensink et al, 1992; Howard and

Wylie-Rosett, 2001; Harvard School of Public Health, 2014) and might adversely promote inflammation and over-activity of the immune system, phenomena that may, it has been posited, be linked to an increased risk of heart disease, stroke, diabetes and other chronic conditions (Mozaffarian et al, 2004; Brown et al, 2010; Iwata et al, 2011). Consumption of trans fat has also been linked to a reduction in the normal healthy responsiveness of endothelial cells (Katan et al, 1995). In addition, a study in primates showed that monkeys fed trans fatty acids gained weight and deposited more central fat even without increased food consumption, and they also displayed impaired glucose tolerance (Kavanagh et al, 2007).

Effect of chemical additives on the body

Practically, it is very difficult to isolate the impact of any one chemical additive on the body. Of relevance to the theme of this piece, though, it has been postulated that among the health risks chemical additives may pose, the storage of such substances in newly created fat cells may impede weight loss (Richards, 2012).

“The imbalance between our current lifestyles and that of our ancestors manifests itself, in part, as an increased risk of obesity and type 2 diabetes, and of ill health in general”



“In offering dietary advice for people with diabetes that will promote weight loss and better health, we need to go beyond simple macronutrient ratios and consider the need to cut back on processed products and instead eat more natural foods.”

Summary

The imbalance between our current lifestyles and that of our ancestors manifests itself, in part, as an increased risk of obesity and type 2 diabetes, and of ill health in general (Leach, 2006; Swedish Council on Health Technology Assessment, 2013). I believe that in offering dietary advice for people with diabetes that will promote weight loss and better health, we need to go beyond simple macronutrient ratios and consider the need to cut back on processed products and, instead, eat more natural foods. ■

Alhazmi A, Stojanovski E, McEvoy M, Garg ML (2012) Macronutrient intakes and development of type 2 diabetes: a systemic review and meta-analysis of cohort studies. *J Am Coll Nutr* **31**: 243–58

Basaranoglu M, Basaranoglu G, Sabuncu T, Sentürk H (2013) Fructose as a key player in the development of fatty liver disease. *World J Gastroenterol* **19**: 1166–72

Bray G, Nielsen S, Popkin B (2004) Consumption of high-fructose corn syrup in beverages may play a role in the epidemic of obesity. *Am J Clin Nutr* **79**: 537–43

Brown MA, Storlien LH, Huang XF et al (2010) Dietary fat and carbohydrate composition: Metabolic disease. In: Montmayeur JP, le Coutre J, eds. *Fat Detection: Taste, Texture, and Post Ingestive Effects*. CRC Press, Boca Raton, FL, USA

Coffey DS (2001) Similarities of prostate and breast cancer: Evolution, diet, and estrogens. *Urology* **57**(Suppl 4A): 31–8

Fallon S, Enig M (2001) *Nourishing Traditions: The Cookbook That Challenges Politically Correct Nutrition and the Diet Dictocrats*. NewTrends, Washington, DC, USA

Goran MI, Ulijaszek SJ, Ventura EE (2013) High fructose corn syrup and diabetes prevalence: a global perspective. *Glob Public Health* **8**: 55–64

Harvard School of Public Health (2014) *Shining the Spotlight on Trans Fats*. Harvard School of Public Health, Boston, MA, USA. Available at: <http://www.hsph.harvard.edu/nutritionsource/transfats/> (accessed 12.03.14)

Hines M (2009) *Human Evolution Diet and Health: The Case for Palaeolithic Nutrition*. Healthy Body Publishing (<http://hbpublishing.com/>) [accessed 12.03.14]

Hockett B, Haws J (2003) Nutritional ecology and diachronic trends in Palaeolithic diet and health. *Evol Anthropol* **12**: 211–6

Howard B, Wylie-Rosett J (2001) Sugar and Cardiovascular Disease. A statement for healthcare professionals from the committee on nutrition of the council on nutrition, physical activity, and metabolism of the American Heart Association. *J Nutr* **131**: 132–46

Iwata NG, Pham M, Rizzo NO et al (2011) Trans fatty acids induce vascular inflammation and reduce vascular nitric oxide production in endothelial cells. *PLoS One* **6**: e29600

Katan MB, Zock PL, Mensink RP (1995) Trans fatty acids and their effects on lipoproteins in humans. *Annu Rev Nutr* **15**: 473–93

Kavanagh K, Jones KL, Sawyer J et al (2007) Trans fat diet induces abdominal obesity and changes in insulin sensitivity in monkeys. *Obesity (Silver Spring)* **15**: 1675–84

Lakhan S, Kirchgessner A (2013) The emerging role of dietary fructose in obesity and cognitive decline. *Nutr J* **12**: 114

Lankinen M, Schwab U, Kolehmainen M (2011) Whole grain products, fish and bilberries alter glucose and lipid metabolism in a randomized, controlled trial: the Sysdimet study. *PLoS One* **6**: e22646

Leach JD, Gibson GR, Van Loo J (2006) Human evolution, nutritional ecology and prebiotics in ancient diet. *Bioscience Microflora* **25**: 1–8

Mack S (2001) *Re: How many cells does a human lose every second?* Available at: <http://www.madsci.org/posts/archives/2001-02/981770369.An.r.html> (accessed 12.03.14)

McMichael AJ (2005) Integrating nutrition with ecology: balancing the health of humans and biosphere. *Public Health Nutr* **8**: 706–15

Mensink RP, Zock PL, Katan MB, Hornstra G (1992) Effect of dietary cis and trans fatty acids on serum lipoprotein[a] levels in humans. *J Lipid Res* **33**: 1493–501

Mozaffarian D, Pischon T, Hankinson SE et al (2004) Dietary intake of trans fatty acids and systemic inflammation in women. *Am J Clin Nutr* **79**: 606–12

NHS Choices (2013) *Do processed baby foods 'lack nutritional value'?* Available at: <http://1.usa.gov/1eacgBA> (accessed 12.03.14)

NutritionData.com (2013) *Nutritional Effects of Food Processing*. Available at: <http://nutritiondata.self.com/topics/processing> (accessed 12.03.14)

Price WA (2003) *Nutrition and Physical Degeneration* (6th edition). Keats Pub, Wilkes-Barre, PA, USA

Richards BJ (2012) *Why Toxins and Waste Products Impede Weight Loss – The Leptin Diet Weight Loss Challenge #3*. Wellness Resources, Minneapolis, MN, USA. Available at: http://www.wellnessresources.com/weight/articles/why_toxins_and_waste_products_impede_weight_loss_-_the_leptin_diet_weight_loss/ (accessed 12.03.14)

Samaniego-Vaesken M, Alonso-Aperte E, Varela-Moreiras G (2012) Vitamin food fortification today. *Food Nutr Res* **56**: 10.3402

Schmidt CW (2005) Genetically modified foods: Breeding uncertainty. *Environ Health Perspect* **113**: A526–A533

Swedish Council on Health Technology Assessment (2013) *Dietary treatment of obesity – A systematic review*. Swedish Council on Health Technology Assessment, Stockholm, Sweden. Available at: http://www.sbu.se/upload/Publikationer/Content1/1/Diets_among_obese_individuals.pdf (accessed 12.03.14)

Van Gysel M, Cossey V, Fieueus S, Schuermans A (2012) Impact of pasteurization on the antibacterial properties of human milk. *Eur J Pediatr* **171**: 1231–7

Webster JL, Dunford EK, Neal BN (2010) A systematic survey of the sodium contents of processed foods. *Am J Clin Nutr* **91**: 413–20